

Product datasheet for **SC336654**

VPS45A (VPS45) (NM_001279354) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	VPS45A (VPS45) (NM_001279354) Human Untagged Clone
Tag:	Tag Free
Symbol:	VPS45
Synonyms:	H1; H1VPS45; SCN5; VPS45A; VPS45B; VPS54A; VSP45; VSP45A
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >SC336654 representing NM_001279354.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)


```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCCGCATCGCC
ATGGTATACACACAATCGGAGATTCTACAGAAGGAAGTGTACCTCTTTGAACGCATTGATTCTCAAAT
CGAGAGATCATGAAACACCTGAAGCAATTTGTTTTCTCGACCTACAAAGGAGAATGTGGATTATATT
ATTCAGGAGCTCCGAAGACCCAAATACACTATATATTTTCAATTTTTCAGTAATGTGATCAGCAAGAGT
GACGTGAAGTCATTGGCTGAAGCTGATGAACAGGAAGTTGTGGCTGAGGTTCCAGGAATTTTATGGTGAT
TACATTGCTGTGAACCCACATTTGTTTTCCCTCAATATTTTGGGTTGCTGCCAGGGTCGAAATTTGGGAT
CCAGCCCAGCTATCTAGAACAACCAAGGGCTTACAGCTCTCCTTTTATCTCTGAAGAAGTGTCCCATG
ATTCGTTATCAGCTCTCATCAGAGGCAGCAAAGAGACTTCGAGAGTGCCTAAGCAAGTGATAACTAAA
GAATATGAACTGTTTGAATTCGTCGGACAGAGGTTCCCTCATTGCTCCTATTTTAGATCGCTGTGAT
GATGCCATCACCCATTGCTAAACCAGTGGACATATCAGGCCATGGTCCACGAACACTAGGCATAAAC
AACAAATCGGATTGATCTTCCAGAGTGCCGGGAATCAGTAAAGACTTAAGAGAAGTGGTCCATCTGCT
GAAAATGATGAATTCTATGCTAATAATGTACCTGAACCTTGGCTGAGATTGGTAGCAATATAAAGAAAT
CTCATGGAAGATTTTCAGAAGAAGAAACAAAAGAACAGCAAAAAGTCTAGAAATCAATAGCAGACATGAAG
GCGTTTGTGAGAATTATCCACAGTTCAAGAAAATGTCTGGGACTGTTTCAAAGCATGTGACAGTGGTT
GGAGAAGTGTCTCGATTGGTCAGTGAACCGAATCTGCTGGAGGTTTCAGAGGTTGAGCAAGAAGTGGCC
TGTCAAATGACCATTCTAGTGTCTCCAGAATATAAAAAGGCTTCTGCAGAACCCCAAAGTGACAGAG
TTTGATGCTGCCCGCTGGTGATGCTTTATGCTTTACATTATGAGCGACACAGCAGCAATAGCCTGCCA
GGACTAATGATGGACCTCAGGAATAAAGGTGTTTCTGAGAAGTATCGAAAGCTCGTGTCTGCAGTTGTT
GAATATGGTGGTAAACGAGTCAGAGGAAGTGACCTTTCAGCCCCAAAGATGCTGTGGCTATCACCAAA
CAATTCCTCAAAGGACTGAAGGGAGTAGAAAATGTATATACACAGCATCAACCTTTCCTACATGAAACC
CTGGATCATCTCAAAAGGAAGGCTTAAAGAAAACCTATATCCTTATTTAGGCCCCAGCACACTCAGA
GACAGACCTCAGGATATCATTGTGTTTGAATTGGAGGAGCCACCTATGAAGAGGCTCTAACAGTTTAT
AACCTGAACCGCACCCTCCTGGAGTGAGGATTGTCTGGGAGGCACCACAGTGCACAACACGAAAAGT
TTCCTAGAGGAAGTTCTGGCTTCTGGACTGCACAGCCGAAGCAAGGAGAGCTCTCAAGTCACATCAAGG
TCAGCGAGCAGAAGATGA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGCGC
  
```

Restriction Sites:

Sgfl-MluI

Plasmid Map:



ACCN:	NM_001279354
Insert Size:	1605 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001279354.1
RefSeq Size:	2469 bp
RefSeq ORF:	1605 bp
Locus ID:	11311
UniProt ID:	Q9NRW7
Cytogenetics:	1q21.2
Protein Pathways:	Endocytosis
MW:	61.1 kDa
Gene Summary:	<p>Vesicle mediated protein sorting plays an important role in segregation of intracellular molecules into distinct organelles. Genetic studies in yeast have identified more than 40 vacuolar protein sorting (VPS) genes involved in vesicle transport to vacuoles. This gene is a member of the Sec1 domain family, and shows a high degree of sequence similarity to mouse, rat and yeast Vps45. The exact function of this gene is not known, but its high expression in peripheral blood mononuclear cells suggests a role in trafficking proteins, including inflammatory mediators. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2013]</p> <p>Transcript Variant: This variant (3) has an alternate splice site in the 5' region, which results in a downstream translation start codon, compared to variant 1. The resulting isoform (3) has a shorter N-terminus, compared to isoform 1.</p>