

## Product datasheet for **SC336849**

### TACC2 (NM\_001291879) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	TACC2 (NM_001291879) Human Untagged Clone
Tag:	Tag Free
Symbol:	TACC2
Synonyms:	AZU-1; ECTACC
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)




[View online »](#)

Fully Sequenced ORF: >SC336849 representing NM\_001291879.  
Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC
ATGCCCTGAGGAGGCCAAGATGAAAAAGACACCCGAGAAACTTGACAACACTCCTGCCTCACCTCCC
AGATCCCCTGCTGAACCAATGACATCCCCATTGCTAAAGGTACTTACACCTTTGATATTGACAAGTGG
GATGACCCCAATTTAAACCCTTTTTCTCCACCTCAAAAATGCAGGAGTCTCCAAAAGTCCCAACAA
TCATACACTTTGACCCAGACACCTGTGATGAGTCCGTTGACCCCTTTAAGACATCCTCTAAGACCCCC
AGCTCACCTTCTAAATCCCCAGCCTCCTTTGAGATCCCAGCCAGTGTATGGAAGCCAATGGAGTGGAC
GGGGATGGGCTAAACAAGCCCGCAAGAAGAAGACGCCCTAAAGACTGACACATTTAGGGTAAAA
AAGTCGCAAAACGGTCTCCTCTCTGATCCACCTTCCAGGACCCACCCAGCTGCTACACCAGAA
ACACCACAGTGTCTCTGCGGTGGTCCACGCCACAGATGAGGAAAAGCTGGCGGTACCAACCAGAAG
TGGACGTGCATGACAGTGGACCTAGAGGCTGACAAACAGGACTACCCGCAGCCCTCGGACCTGTCCACC
TTTGTAAACGAGACCAAAATTCAGTTCACCCACTGAGGAGTTGGATTACAGAACTCCTATGAAATTGAA
TATATGGAGAAAATTGGCTCCTCTACCTCAGGACGACGATGCCCCGAAGAAGCAGGCCTTGACCTT
ATGTTTGACACTTCTCAGGAGAGCCCTGTCAAGTCATCTCCCGTCCGCATGTGAGAGTCCCCGACGCCG
TGTTTCAGGGTCAAGTTTTGAAGAGACTGAAGCCCTTGTGAACACTGCTGCGAAAAACCAGCATCCTGTC
CCACGAGGACTGGCCCTAACCAAGAGTCACACTTGCAGGTGCCAGAGAAATCCTCCAGAAGGAGCTG
GAGGCCATGGGCTTGGGCACCCCTTCAAGCGATTGAAATTAGAGAGGCTGCTCACCAACAGACGTC
TCCATCTCAAAAACAGCCTTGTACTCCCGCATCGGGACCGCTGAGGTGGAGAACTGCAGGCCTTCTG
TTCCAGCAGCCCGACCTGGACTCTGCCCTCCAGATCGCCAGAGCAGAGATCATAACCAAGGAGAGAGAG
GTCTCAGAAATGGAAAGATAAATATGAAGAAAGCAGGCGGGAAGTATGGAATGAGGAAAAATAGTGCC
GAGTATGAGAAGACCATCGCTCAGATGATAGAGGACGAACAGAGAGAGAAGTCAGTCTCCACAGACG
GTGCAGCAGCTGGTTCTGGAGAAGGAGCAAGCCCTGGCCGACCTGAACTCCGTGGAGAAGTCTCTGGCC
GACCTCTCAGAAGATATGAGAAGATGAAGGAGGTCTAGAAGGCTTCCGCAAGAATGAAGAGGTGTTG
AAGAGATGTGCGCAGGAGTACCTGTCCCGGTGAAGAAGGAGGAGCAGAGGTACCAGGCCCTGAAGGTG
CACGCGGAGGAGAACTGGACAGGGCAATGCTGAGATTGCTCAGGTTGAGGCAAGGCCAGCAGGAG
CAAGCCGCCACCAGGCCAGCCTGCGGAAGGAGCAGCTGCGAGTGGACGCCCTGGAAAGGACGCTGGAG
CAGAAGAATAAAGAAATAGAAGAACTACCAAGATTTGTGACGAAGTATTGCCAAAATGGGGAAAAGC
TAA
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTAAACGGCCGGC
```

Restriction Sites: SgfI-MluI

**Plasmid Map:**


**ACCN:** NM\_001291879

**Insert Size:** 1728 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_001291879.1](#)

**RefSeq Size:** 3709 bp

**RefSeq ORF:** 1728 bp

**Locus ID:** 10579

**Cytogenetics:** 10q26.13

**MW:** 64.7 kDa

**Gene Summary:**

Transforming acidic coiled-coil proteins are a conserved family of centrosome- and microtubule-interacting proteins that are implicated in cancer. This gene encodes a protein that concentrates at centrosomes throughout the cell cycle. This gene lies within a chromosomal region associated with tumorigenesis. Expression of this gene is induced by erythropoietin and is thought to affect the progression of breast tumors. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (8) differs in the 5' UTR and coding region and lacks two internal coding exons compared to variant 1. These differences result in the use of a downstream AUG compared to variant 1. The encoded isoform (h) has a shorter N-terminus compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.