

## Product datasheet for **SC336925**

### NR2C2 (NM\_001291694) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	NR2C2 (NM_001291694) Human Untagged Clone
Tag:	Tag Free
Symbol:	NR2C2
Synonyms:	TAK1; TR4
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)




[View online »](#)

Fully Sequenced ORF: >SC336925 representing NM\_001291694.  
Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCCGCATCGCC
ATGACCAGCCCCCCCCACGCATCCAGATAATCTCCACCGACTCTGCTGTAGCCTCACCTCAGCGCATT
CAGATTGTACAGACCAGCAGACAGGACAGAAAAATCCAGATAGTCACCGCAGTGGACGCCTCCGGATCC
CCCAAACAGCAGTTCATCCTGACCAGCCCAGATGGAGCTGGAACCTGGGAAGGTGATCCTGGCTTCCCA
GAGACATCCAGCGCCAAGCAACTCATATTCACCACCTCAGACAACCTCGTCCCTGGCAGGATCCAGATT
GTCACGGATTCTGCCTCTGTGGAGCGTTTACTGGGAAGACGGACGTCCAGCGGCCCCAGGTGGTAGAG
TACTGTGTGGTCTGTGGCGACAAAGCCTCCGGCCGTCCTATGGGGCTGTGAGTTGTGAAGTTGCAAA
GGTTTCTTCAAAGGAGTGTGAGGAAAAATTTGACCTACAGCTGCCGGAGCAACCAAGACTGCATCATC
AATAAACATCACCGAACCGCTGTGAGTTTGGCCGCTGAAAAATGCTTAGAGATGGGCATGAAAAATG
GAATCTGTGCAGAGTGAACGGAACCGCTTCGATGTGCAACGGGAGAAACCAAGCAATTGTGCTGCTCA
ACTGAGAAAAATCTATATCCGAAAGACCTGAGAAGTCCCCTGATAGCTACTCCCAGCTTTGTGGCAGAC
AAAGATGGAGCAAGACAACAGGCTTCTTGTATCCAGGGATGCTTGTGAACATCCAGCAGCTTTGATA
CGTGAGGATGGTACAGTTCTCCTGGCCACGGATTCTAAGGCTGAAACAAGCCAGGAGCTCTGGGCACA
CTGGCAAATGTAGTGACCTCCCTTGCCAACTAAGTGAATCTTTGAAACAAGGTGACACTTCAGAAATC
CAGCCAGAGGACCAGTCTGCAAGTGAGATAACTCGGGCATTGATACCTTAGCTAAAGCACTTAATACC
ACAGACAGCTCCTCTTCTCCAAGCTTGGCAGATGGGATAGACACCAGTGGAGGAGGGAGCATCCACGTC
ATCAGCAGAGACCAGTCGACACCCATCATTGAGGTTGAAGGCCCTCCTTTGAGACACACAGTCACA
TTTAAGCTAACAATGCCAGTCCAATGCCAGAGTACCTCAACGTGCACTACATCTGTGAGTCTGCATCC
CGTCTGCTTTTCTCTCAATGCACTGGGCTCGGTCAATCCAGCCTTTTCAGGCAGTTGGCAGGACTGC
AACACCAGCCTTGTGCGGCCTGCTGGAATGAGCTCTTACCCTCGGCCTGGCCAGTGTGCCAGGTC
ATGAGTCTCTCCACCATCCTGGCTGCCATTGTCAACCACCTGCAGAACAGCATCCAGGAAGATAAATT
TCTGGTGACCGGATAAAGCAAGTCATGGAGCACATCTGGAAGCTGCAGGAGTTCTGTAACAGCATGGCG
AAGCTGGATATAGATGGCTATGAGTATGCATACCTTAAAGCTATAGTTCTCTTTAGCCCCGATCATCCA
GGTTTGACCAGCACAAGCCAGATTGAAAAATCCAAGAAAAGGCACAGATGGAGTTGCAGGACTATGTT
CAGAAAACCTACTCAGAAGACACCTACCGATTGGCCCGGATCCTCGTTCGCCTGCCGGCACTCAGGCTG
ATGAGCTCCAACATAACAGAAGAACTTTTTTTTACTGGTCTCATTGGCAATGTTTCGATAGACAGCATA
ATCCCCTACATCCTCAAGATGGAGACAGCAGAGTATAATGGCCAGATCACCGGAGCCAGTCTATAG
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
```

Restriction Sites: SgfI-MluI

**Plasmid Map:**


**ACCN:** NM\_001291694

**Insert Size:** 1791 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_001291694.1](#)

**RefSeq Size:** 8428 bp

**RefSeq ORF:** 1791 bp

**Locus ID:** 7182

**UniProt ID:** [P49116](#)

**Cytogenetics:** 3p25.1

**Protein Families:** Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

**MW:** 65.4 kDa

**Gene Summary:**

This gene encodes a protein that belongs to the nuclear hormone receptor family. Members of this family act as ligand-activated transcription factors and function in many biological processes such as development, cellular differentiation and homeostasis. The activated receptor/ligand complex is translocated to the nucleus where it binds to hormone response elements of target genes. The protein encoded by this gene plays a role in protecting cells from oxidative stress and damage induced by ionizing radiation. The lack of a similar gene in mouse results in growth retardation, severe spinal curvature, subfertility, premature aging, and prostatic intraepithelial neoplasia (PIN) development. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Apr 2014]

**Transcript Variant:** This variant (2) lacks an in-frame exon in the 5' coding region compared to variant 1. The encoded isoform (2) is shorter than isoform 1.

**Sequence Note:** This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.