

Product datasheet for **SC336928**

LILRB2 (NM_001278403) Human Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: LILRB2 (NM_001278403) Human Untagged Clone
Tag: Tag Free
Symbol: LILRB2
Synonyms: CD85D; ILT-4; ILT4; LIR-2; LIR2; MIR-10; MIR10
Vector: pCMV6-Entry (PS100001)
Fully Sequenced ORF: >SC336928 representing NM_001278403.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```


ATGACCCCATCGTCACAGTCTGATCTGTCTCGGGCTGAGTCTGGGCCCCAGGACCCGCGTGCAGACA
GGGACCATCCCCAAGCCACCCTGTGGGCTGAGCCAGACTCTGTGATCAGCCAGGGGAGTCCCGTACC
CTCAGTTGTGAGGGGAGCCTTGAAGCCAGGAGTACCGTCTATATAGGGAGAAAAATCAGCATCTTGG
ATTACACGGATACGACCAGAGCTTGTGAAGAACGGCCAGTCCACATCCCATCCATCACCTGGGAACAC
ACAGGGCGATATGGCTGTCAGTATTACAGCCGCGCTCGGTGGTCTGAGCTCAGTACCCCTGGTGTG
GTGATGACAGGAGCCTACCCAAAACCCACCTCTCAGCCAGCCAGCCCTGTGGTGACCTCAGGAGGA
AGGGTGACCTCCAGTGTGAGTACAGGTGGCATTGGCGGCTTATTCTGTGTAAAGGAAGGAGAAGAT
GAACACCCACAATGCCTGAACTCCAGCCCATGCCCCGTGGGTGCTCCCGGCCATCTTCTCCGTGGGC
CCCGTGAGCCCAATCGCAGGTGGTGCACAGGTGCTATGGTTATGACTTGAAGTCTCCCTATGTGTGG
TCTTACCCAGTGTCTCCTGGAGCTCCTGGTCCCAGGTGTTTCTAAGAAGCCATCACTCTCAGTGCAG
CCGGTCTGTGATGGCCCTGGGGAAAGCCTGACCTCCAGTGTGTCTCTGATGTCGGCTATGACAGA
TTTGTCTGTACAAGGAGGGGAAACGTGACCTTCGCCAGTCCCTGGCCGGCAGCCCAAGGCTGGGCTC
TCCAGGCCAACTTACCCTGGGCCCTGTGAGCCGCTCCTACGGGGCCAGTACAGATGCTACGGTGCA
CACAACCTCTCCTGAGTGTGCGCCCCAGCAGCCCTGGACATCCTGATCACAGGACAGATCCGT
GGCACACCCTTCACTCAGTGCAGCCAGGCCCCACAGTGGCCTCAGGAGAGAAGCTGACCTGCTGTGT
CAGTCATGGCGGAGTCCACACTTTCCTTCTGACCAAGGCGGGAGCAGTGTGCCCCACTCCGCTCTA
AGATCAATACACGAATATCCTAAGTACCAGGCTGAATTCCCCATGAGTCTGTGACCTCAGCCACGCG
GGGACCTACAGGTGCTACGGTCACTCACTCCGACCCCTACCTGCTGTCTACCCCAAGTGGCCCTG
GAGCTCGTGGTCTCAGGACCTCCATGGTTCAGCCCCCACCACCGTCCCCTCCACACCTGGC
CCTGAGGACCAGCCCTACCCCACTGGGTGGATCCCCAAAGTGGTCTGGGAAGGCACCTGGGGTT
GTGATCGGCATCTTGGTGGCCGTGCTCTACTGCTCCTCCTCCTCCTCCTCCTCCTCCTCCTCCTCGA
CATCGACGTGAGGCAAACTGGACATCGACCCAGAGAAAGGCTGATTTCCAACATCCTGCAGGGGCT
GTGGGGCCAGAGCCACAGACAGAGGCCTGCAGTGGAGTCCAGCCAGCTGCCGACGCCAGGAAGAA
AACCTCTATGCTGCCGTGAAGGACACACAGCCTGAAGATGGGGTGGAGATGGACACTCGGCTGCTGCA
TCTGAAGCCCCCAGGATGTGACCTACGCCAGCTGCACAGCTTGACCCTCAGACGGAAGGCAACTGAG
CCTCCTCCATCCCAGGAAAGGGAACCTCCAGCTGAGCCAGCATCTACGCCACCTGGCCATCCACTAG
  
```



[View online »](#)

Restriction Sites: Sgfl-MluI

Plasmid Map:



ACCN: NM_001278403

Insert Size: 1794 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001278403.2](#)

RefSeq Size: 2799 bp

RefSeq ORF: 1794 bp

Locus ID: 10288

Cytogenetics: 19q13.42

Protein Families: Transmembrane

MW: 65 kDa

Gene Summary:

This gene is a member of the leukocyte immunoglobulin-like receptor (LIR) family, which is found in a gene cluster at chromosomal region 19q13.4. The encoded protein belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The receptor is expressed on immune cells where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (3) differs in the 5' UTR and uses an alternate in-frame splice site in the central coding region, compared to variant 1. The encoded isoform (2) is shorter, compared to isoform 1. Both variants 2 and 3 encode the same isoform.