

# **Product datasheet for SC336985**

## BRSK2 (NM\_001282218) Human Untagged Clone

### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** BRSK2 (NM\_001282218) Human Untagged Clone

Tag: Tag Free Symbol: BRSK2

Synonyms: C11orf7; PEN11B; SAD1; SADA; STK29

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn





**Fully Sequenced ORF:** 

>SC336985 representing NM\_001282218.

Blue=Insert sequence Red=Cloning site Green=Tag(s)


GCTCGTTTAGTGAACCGTCAGAATTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCCGCCGCGGATCGCC

ATGAAGGTGGAGCGGGAGATCGCGATCCTGAAGCTCATTGAGCACCCCCACGTCCTAAAGCTGCACGAC GTTTATGAAAACAAAAATATTTGTACCTGGTGCTAGAACACGTGTCAGGTGGTGAGCTCTTCGACTAC CTGGTGAAGAAGGGGAGGCTGACGCCTAAGGAGGCTCGGAAGTTCTTCCGGCAGATCATCTCTGCGCTG GACTTCTGCCACAGCCACTCCATATGCCACAGGGATCTGAAAACCTGAAAACCTCCTGCTGGACGAGAAG AACAACATCCGCATCGCAGACTTTGGCATGGCGTCCCTGCAGGTTGGCGACAGCCTGTTGGAGACCAGC TGTGGGTCCCCCACTACGCCTGCCCCGAGGTGATCCGGGGGGGAGAAGTATGACGGCCGGAAGGCGGAC GTGTGGAGCTGCGGCGTCATCCTGTTCGCCTTGCTGGTGGGGGCTCTGCCCTTCGACGATGACAACTTG AGTCTGCTACGGGGCATGATCGAGGTGGACGCCGCACGCCCTCACGCTAGAGCACATTCAGAAACAC ATATGGTATATAGGGGGCAAGAATGAGCCCGAACCAGAGCAGCCCATTCCTCGCAAGGTGCAGATCCGC GACCGCAACAAGCTGCTGCAGGACCTGCTGTCCGAGGAGGAGCACCAGGAGAAGATGATTTACTTCCTC CTCCTGGACCGGAAAGAAAGGTACCCGAGCCAGGAGGATGAGGACCTGCCCCCCGGAACGAGATAGAC CCTCCCGGAAGCGTGTGGACTCCCCGATGCTGAACCGGCACGGCAAGCGGCCAGAACGCAAATCC ATGGAGGTGCTCAGCGTGACGGACGGCGGCTCCCCGGTGCCTGCGCGGGGGCCATTGAGATGGCCCAG CACGGCCAGAGGTCTCGGTCCATCAGCGGTGCCTCCTCAGGCCTTTCCACCAGCCCACTCAGCAGCCCC CGGGTGACCCCTCACCCTCACCAAGGGGCAGTCCCCTCCCCACCCCCAAGGGGACACCTGTCCACACG CCAAAGGAGAGCCCGGCTGGCACGCCCAACCCCACGCCCCGTCCAGCCCCAGCGTCGGAGGGGTGCCC TGGAGGGCGCGCTCAACTCCATCAAGAACAGCTTTCTGGGCTCACCCCGCTTCCACCGCCGGAAACTG CAAGTTCCGACGCCGGAGGAGATGTCCAACCTGACACCAGAGTCGTCCCCAGAGCTGGCGAAGAAGTCC TGGTTTGGGAACTTCATCAGCCTGGAGAAGGAGGAGCAGATCTTCGTGGTCATCAAAGACAAACCTCTG AGCTCCATCAAGGCTGACATCGTGCACGCCTTCCTGTCGATTCCCAGTCTCAGCCACAGCGTCATCTCC CAAACGAGCTTCCGGGCCGAGTACAAGGCCACGGGGGGGCCCAGCCGTGTTCCAGAAGCCGGTCAAGTTC CAGGTTGATATCACCTACACGGAGGGTGGGGAGGCGCAGAAGGAGAACGGCATCTACTCCGTCACCTTC ACCCTGCTCTCAGGCCCCAGCCGTCGCTTCAAGAGGGTGGTGGAGACCATCCAGGCCCAGCTGCTGAGC ACACACGACCCGCCTGCGGCCCAGCACTTGTCAGAACCCCCCCACCAGCGCCCAGGACTAAGCTGGGGT GCTGGGCTTAAGGGCCAGAAGGTGGCCACCAGCTACGAGAGTAGCCTCTGA

**Restriction Sites:** Sgfl-Mlul



#### Plasmid Map:



**ACCN:** NM\_001282218

**Insert Size:** 1845 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001282218.1</u>

RefSeq Size: 4127 bp RefSeq ORF: 1845 bp Locus ID: 9024

UniProt ID: Q8IWQ3

Cytogenetics: 11p15.5

**Protein Families:** Druggable Genome, Protein Kinase

MW: 68.7 kDa



#### **Gene Summary:**

Serine/threonine-protein kinase that plays a key role in polarization of neurons and axonogenesis, cell cycle progress and insulin secretion. Phosphorylates CDK16, CDC25C, MAPT/TAU, PAK1 and WEE1. Following phosphorylation and activation by STK11/LKB1, acts as a key regulator of polarization of cortical neurons, probably by mediating phosphorylation of microtubule-associated proteins such as MAPT/TAU at 'Thr-529' and 'Ser-579'. Also regulates neuron polarization by mediating phosphorylation of WEE1 at 'Ser-642' in postmitotic neurons, leading to down-regulate WEE1 activity in polarized neurons. Plays a role in the regulation of the mitotic cell cycle progress and the onset of mitosis. Plays a role in the regulation of insulin secretion in response to elevated glucose levels, probably via phosphorylation of CDK16 and PAK1. While BRSK2 phosphorylated at Thr-174 can inhibit insulin secretion (PubMed:22798068), BRSK2 phosphorylated at Thr-260 can promote insulin secretion (PubMed:22669945). Regulates reorganization of the actin cytoskeleton. May play a role in the apoptotic response triggered by endoplasmic reticulum (ER) stress. [UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (6) differs in both UTRs and has multiple differences in the coding region, one of which results in a translational frameshift, compared to variant 1. The resulting protein (isoform 5) has a shorter N-terminus and a distinct C-terminus compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.