

Product datasheet for **SC337250**

IL12RB1 (NM_001290024) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	IL12RB1 (NM_001290024) Human Untagged Clone
Tag:	Tag Free
Symbol:	IL12RB1
Synonyms:	CD212; IL-12R-BETA1; IL12RB; IMD30
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001290024, the custom clone sequence may differ by one or more nucleotides

```

ATGTTTCGTTGGTCTTTTCTCCTTGCTCAGCTTCAATGTGTTCCGGAGTGGGGACGGGGTGGCTGAACCTC
GCAGGTGGCAGAGAGGCTCCCTGGGGCTGTGGGGCTCTACGTGGATCCGATGGAGCCGCTGGTGACCTG
GGTGGTCCCCCTCCTCTTCTCCTTCTCCTGTGTCCAGGCAGGGCGCTGCCTGCAGAACCAGTGAGTGCTGT
TTTCAGGACCCGCATATCCGGATGCAGACTCAGGCTCGGCCCTCGGGCCCTAGGGACCTGAGATGCTATC
GGATATCCAGTGATCGTTACGAGTGCTCCTGGCAGTATGAGGGTCCCACAGCTGGGGTACGCCACTTCTCT
GCGGTGTTGCCTTAGCTCCGGGGCTGTGCTACTTCCGCCCGGCTCAGCCACCAGGCTGCAGTCTCTCC
GACCAGGCTGGGGTGTCTGTGCTGTACACTGTCACACTCTGGGTGGAATCCTGGGCCAGGAACCAGACAG
AGAAGTCTCCTGAGGTGACCCTGCAGCTCTACAACCTCAGTTAAATATGAGCCTCCTCTGGGAGACATCAA
GGTGTCCAAGTTGGCCGGGCAGCTGCGTATGGAGTGGGAGACCCCGGATAACCAGTTGGTGTGAGGTG
CAGTTCGGCACCCGACCCAGCAGCCCATGGAAGTTGGGCGACTGCGGACCTCAGGATGATGATACTG
AGTCTGCTCTGCCCTGGAGATGAATGTGGCCAGGAATTCAGCTCCGACGACGGCAGCTGGGGAG
CCAAGGAAGTTCCTGGAGCAAGTGGAGCAGCCCGTGTGCGTTCCCCCTGAAAACCCCCACAGCCTCAG
GTGAGATTCTCGGTGGAGCAGCTGGGCCAGGATGGGAGGAGGCGGCTGACCCTGAAAGAGCAGCCAAACC
AGCTGGAGCTTCCAGAAGGCTGTCAAGGGCTGGCGCTGGCAGGAGTCACTTACCAGTACAGCTCCA
CATGCTGTCTGCCCGTGTAAAGGCCAAGGCCACCAGGACCCTGCACCTGGGGAAGATGCCCTATCTCTCG
GGTGTGCCTACAACGTGGCTGTCTCTCCTCGAACCAATTTGGTCTGGCCTGAACCAGACGTGGCACA
TTCTGCCGACACCCACAGAACCAAGTGGCTCTGAATATCAGCGTCGGAACCAACGGGACCACCATGTA
TTGGCCAGCCCGGGCTCAGAGCATGACGTATTGCATTGAATGGCAGCCTGTGGCCAGGACGGGGCCTT
GCCACCTGCAGCCTGACTGCGCCGCAAGACCCGGATCCGGCTGGAATGGCAACCTACAGTGGAGTCCGAG
AGTCTGGGGCAATGGGGCAGGAAAAGTGTACTACATTACCATCTTTGCCCTCGCGACCCCGAGAAGCT
CACCTTGTGGTCTACGGTCTGTCCACCTACCCTTTGGGGCAATGCCTCAGCAGCTGGGACACCCGCAC
CACGTCTCGGTGAAGAATCATAGCTTGGACTCTGTGTCTGTGGACTGGGCACCATCCCTGCTGAGCACCT
GTCCCGCGCTCTAAAGGAGTATGTTGTCCGCTGCCGAGATGAAGACAGCAAACAGGTGTGAGAGCATCC
CGTGCAGCCACAGAGACCCAAGTTACCCTCAGTGGCCTGCGGGCTGGTGTAGCCTACACGGTGCAGGTG
CGAGCAGACACAGCGTGGCTGAGGGGTGTCTGGAGCCAGCCCAGCGCTTACGATCGAAGTGCAGGTTT
CTGATTGGCTCATCTTCTCGCCTCCCTGGGGAGCTTCTGAGCATCCTTCTCGTGGGCGTCTTGGCTA
CCTTGGCCTGAACAGGGCCGACGGCACCTGTGCCCGCCGCTGCCACACCTGTGCCAGCTCCGCCATT
GAGTTCCTGGAGGGAAGGAGACTTGGCAGTGGATCAACCCAGTGGACTTCCAGGAAGAGGCATCCCTGC
AGGAGGCCCTGGTGGTAGAGATGCTCTGGGACAAAGGCGAGAGGACTGAGCCTCTCGAGAAGACAGAGCT
ACCTGAGGGTGCCCTGAGCTGGCCCTGGATACAGAGTTGTCCTTGGAGGATGGAGACAGGTGCAAGGCC
AAGATGTGA
    
```

Restriction Sites: SgfI-MluI

ACCN: NM_001290024

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001290024.1</u> , <u>NP_001276953.1</u>
RefSeq Size:	2791 bp
RefSeq ORF:	2109 bp
Locus ID:	3594
Cytogenetics:	19p13.11
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway
Gene Summary:	<p>The protein encoded by this gene is a type I transmembrane protein that belongs to the hemopoietin receptor superfamily. This protein binds to interleukine 12 (IL12) with a low affinity, and is thought to be a part of IL12 receptor complex. This protein forms a disulfide-linked oligomer, which is required for its IL12 binding activity. The coexpression of this and IL12RB2 proteins was shown to lead to the formation of high-affinity IL12 binding sites and reconstitution of IL12 dependent signaling. Mutations in this gene impair the development of interleukin-17-producing T lymphocytes and result in increased susceptibility to mycobacterial and Salmonella infections. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2014]</p> <p>Transcript Variant: This variant (4) encodes the longest isoform (4).</p>