

Product datasheet for **SC337324**

BRRN1 (NCAPH) (NM_001281710) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	BRRN1 (NCAPH) (NM_001281710) Human Untagged Clone
Tag:	Tag Free
Symbol:	NCAPH
Synonyms:	BRRN1; CAP-H; CAPH; MCPH23
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001281710, the custom clone sequence may differ by one or more nucleotides

```

ATGGGACCTCCCGGCCAGAGACGGAGGACACCCACAGTGCCTCCTCCTTCAGAGCGTGTGTTCC
CGATGCCCTGCCAGGAAGCGCCTCTCAATATTCCTGGCACCCAGTCTCGAAGACTTTCCTCAGAA
TGACGATGAGAAGGAGCGGCTGCAGCGGAGCGCTCGAGGGTCTTTGATCTGCAGTTCAGCACTGACTCA
CCTCGCTTATTGGCCTCCCTCCAGCAGGAGTATTGACATTTAGCTACTATCCCAAGTTTACAACA
CGCAGATTACGGAACATTACTCCACCTGTATCAAAGTGTCCACTGAAAATAAAATCACTACCAAGATGC
TTTTGGTTGCACTTGATTGATTTATGTGAGAGATTCTTAAACAGAAAGACACCGAACCAACCACTTT
AAAGTGGTGGGTTACTCTGGATGCCAGCACCAAGATCTATGCTGTGCGGTGGATGCCGTCCATGCCG
ATGTATACAGAGTCTTGGGGGCTGGGCAAAGATGCACCGTCTTTGGAAGAAGTAGAAGGCCATGTTGC
TGATGGAAGTGTACTGAAATGGGAACAACCAAAAAGGCTGTAAGCCAAAGAAGAAGCACTTACACAGA
ACTATTGAGCAGAACATAAACAACCTCAATGTCTCCGAAGCAGATCGGAAGTGTGAGATTGATCCCATGT
TTCAGAAGACAGCAGCCTCATTTGATGAGTGCAGCACAGCAGGGGTGTTTCTGTCCACTCTCCACTGCCA
GGACTACAGAAGTGAAGTGTCTTCCCTGTATGTCAGACTCTCTCCACGGGAGAACCTCTCGAGTTG
CCAGAGTTAGGTTGTGTAGAAATGACAGATTTAAAGCGCCCTTGACAGCAGTGTGAGAAGATCGCCAGA
TCTGCCCTTCCCTGGCCGGTCCAGTTTACACAGTGGGACAGTAAACACATAATGAGTCTGTGTCCGC
CCTGGTAGACAAGTTTAAAGAAGATGACCAGGTATTTGACATCAATGCTGAAGTTGACGAGAGTGACTGT
GGAGACTTCCCGATGGGTCCCTGGGGGATGACTTTGATGCCAACGATGAACCTGACCACACCGCAGTTG
GGGATCATGAAGAGTTCAGGAGCTGGAAGGAGCCCTGCCAGGTTGAGAGTGCAGGAAGAAATGATTTT
CCTTGGGGATGGAGACATCAGGACCATGTGCCCTTCTGTCTATGAAACCTGGAGAATATTCTTATTTT
AGTCCCTCGGACCATGTCGATGTGGGCTGGCCCGGATCACTGGCGCTTTAGCCCTCGACGCAACAAGATG
CTCCTTCCAATCAGAAAACAAAAAGAAGTACAAAAAAGATTTTGAATTGACTTTGAAGATGATAT
TGACTTTGATGTATTTTAGAAAAACAAAGGCTGCTACTATTCTGACCAAGTCCACTTTGGAGAACCAG
AATTGGAGAGCTACCACCTTCTACAGATTTCAACTACAATGTTGACTCTGTTCCAGCTTCCACTCA
AACCAGGCACCAGGTTACTTAAGATGGCCAGGCCATAGGGTAGAGACTGAGCATTATGAAGAAATTGA
AGACTATGATTACAACAACCTAACGACACCTCCAACCTTTGCCCTGGATTACAGGCTGCTGACAGTGT
GATGAAGATTTGGATGACTTATTTGTGGACCTGTTGGAACTCTGACCTCTCACCTTATCCTTGCCATC
CACCTAAGACAGCACAACAGAATGGTGACTCCAGAAGCCAAAGGATTAGACATACAACATATGGGGA
GTCAAACCTTGGTAGCTGAGCCTCAGAAGTAAATAAAATTGAAATCACTATGCCAAGACTGCCAAAAAG
ATGGACATGAAGAACTGAAGCAGAGCATGTGGAGTCTGCTGACAGCGCTCTCCGAAAGGAGGCAGATG
CAGAGGCAAAACCAGGGAAGCTGGAAAAGAAGCGGCCCTGGCAGAAGTGGCTGACGAGAAGATGCTTAG
CGGGCTCACGAAGGACCTGCAGAGGAGCCTGCCCTGTGATGGCTCAGAACCTCTCCATACCTCTGGCT
TTTGCCTGTCTCTACATTTAGCCAATGAAAAGAACTAAAACCTGGAAGGAACAGAGGACCTCTCTGATG
TTCTTGTGAGGCAAGGAGATTGA
    
```

- Restriction Sites:** SgfI-MluI
- ACCN:** NM_001281710
- OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
- Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001281710.1](#), [NP_001268639.1](#)

RefSeq Size: 6026 bp

RefSeq ORF: 2193 bp

Locus ID: 23397

UniProt ID: [Q15003](#)

Cytogenetics: 2q11.2

Gene Summary: This gene encodes a member of the barr gene family and a regulatory subunit of the condensin complex. This complex is required for the conversion of interphase chromatin into condensed chromosomes. The protein encoded by this gene is associated with mitotic chromosomes, except during the early phase of chromosome condensation. During interphase, the protein has a distinct punctate nucleolar localization. Alternatively spliced transcript variants encoding different proteins have been described. [provided by RefSeq, Jul 2013]

Transcript Variant: This variant (2) uses an alternate splice site in the 5' coding region compared to variant 1. The resulting protein (isoform 2) is shorter but has the same N- and C-termini compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.