

Product datasheet for **SC337328**

MED15 (NM_001293234) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	MED15 (NM_001293234) Human Untagged Clone
Tag:	Tag Free
Symbol:	MED15
Synonyms:	ARC105; CAG7A; CTG7A; PCQAP; TIG-1; TIG1; TNRC7
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001293234, the custom clone sequence may differ by one or more nucleotides

```

ATGGACGTTTCCGGCAAGAGACCGACTGGCGGAGCACCGCTTCCGGCAGAAGCTGGTCAGTCAAATCG
AGGATGCCATGAGGAAAGCTGGTGTGGCACACAGTAAATCCAGCAAGGATATGGAGAGCCATGTTTTCTC
GAAGGCCAAGACCCGGGACGAATACCTTTCTCTCGTGGCCAGGCTCATTATCCATTTTCGAGACATTCAT
AACAAAGAAATCTCAAGCTTCCGTCAAGTATCCTATGAATGCACTCCAGAGCCTGACTGGCGGACCTGCTG
CGGGAGCCGCTGGAATTGGCATGCCTCCTCGGGGCCCGGACAGTCTCTGGGCGGGATGGGTAGCCTTGG
TGCCATGGGACAGCCAATGTCTCTCTCAGGGCAGCCGCCTCCTGGGACCTCGGGGATGGCCCTCACAGC
ATGGCTGTCTGTCTACGGCAACTCCACAGACCCAGCTGCAGCTCCAGCAGGTGGCGCTGCAGCAGCAGC
AGCAACAGCAGCAGTTCAGCAGCAGCAGCAGGCGGCGCTACAGCAGCAGCAGCAGCAGCAGCAACAGCA
GCAGTTCAGGCTCAGCAGAGTCCATGCAGCAGCAGTCCAAGCAGTAGTGCAGCAGCAGCAGCAGCTC
CAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGC
AGCAACAGCAGCTGCAGCGAATAGCACAGCTGCAGCTCCAACAACAGCAACAGCAGCAGCAGCAGCAGCA
GCAGCAGCAGCAGCAGGCTTTGCAGGCCAGCCACCAATTTCAGCAGCCACCGATGCAGCAGCCACAGCCT
CGCCCTCCCAGGCTCTGCCCCAGCAGCTGCAGCAGATGCATCACACACAGCACCACAGCCGCCACCAC
AGCCCCAGCAGCTCCAGTTGCTCAGAACAACCATCACAACTCCCGCCACAGTCGAGCAGCCAGCCTTT
GGTGTACAGGCGCAAGCTCTCCCTGGACAAATGTTGTATACCCAACCACCAGTAAATTTGTCAGGCT
CCGATGGTGGTGCAGCAGCCCCAGTGCAGCCCCAGGTGCAGCAGCAGCAGCAGCAGCAGTACAGACAGCTC
AGGCTGCCAGATGGTGGCTCCCGAGTCCAGGTGCAGACCCCGAGTCGATGCCCTCCCCCAGCC
GTCCCCGAGCCCGGCCAGCCAGCTCACAGCCCAACTCCAACGTAGCTCTGGCCCTGCCCATCTCCC
AGTAGTTCTGCCCAGCCCTCACCGCAGCCCTCCCAGAGCCAGTGACGGCGGGACCCACAGAACT
TCAGTGTCCCCTCACCTGGACCTTTAAACACACCTGTGAACCCAGCTCTGTGATGAGCCAGCTGGCTC
CAGCCAGGCTGAGGAGCAGCAGTACCTGGACAAGCTGAAGCAGCTGTGCAAGTACATCGAGCCCTGCGC
CGCATGATCAACAAGATCGACAAGAACGAAGACAGAAAAAAGGACCTGAGTAAGATGAAGAGCCTTCTGG
ACATTCTGACAGACCCCTCGAAGCGGTGTCCCCTGAAGACCTTGCAAAAGTGTGAGATCGCCCTGGAGAA
ACTCAAGAATGACATGGCGGTGCCACTCCCCACCGCCCCGGTGCCACCGACCAACAGCAGTACCTA
TGCCAGCCGCTCCTGGATGCCGTCTGGCAACATCCGCTCACCTGTCTCAACCATTCCCTGTACCGCA
CATTCTGTCCAGCCATGACCGCCATTACGGCCCAACCATCACGGCCCCAGTGGTGTGACCCGGAAGCG
CAGGCTTGAGGATGATGAGCGGCAGAGCATCCCCAGTGTGCTCCAGGGTGAGGTGCCAGGCTGGACCCC
AAGTTCCTGGTAAACCTGGACCTTCTCACTGCAGCAACAATGGCACTGTCCACCTGATCTGCAAGCTGG
ATGACAAGGACCTCCCAAGTGTGCCACCACTGGAGCTCAGTGTGCCCGCTGACTATCCTGCCAAAGCCC
GCTGTGGATAGACCGGCAGTGGCAGTACGACGCCAACCCCTTCTCCAGTCGGTGCACCGCTGCATGACC
TCCAGGCTGTGCAGCTCCCGGACAAGCACTCGGTACCAGCTTGTCTAACACCTGGGCCAGAGCGTCC
ACCAGGCTGCCTCTCAGCCGCCTAG
    
```

- Restriction Sites:** SgfI-MluI
- ACCN:** NM_001293234
- OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
- Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001293234.1, NP_001280163.1</u>
RefSeq Size:	3280 bp
RefSeq ORF:	2196 bp
Locus ID:	51586
Cytogenetics:	22q11.21
Protein Families:	Druggable Genome, Transcription Factors
Gene Summary:	<p>The protein encoded by this gene is a subunit of the multiprotein complexes PC2 and ARC/DRIP and may function as a transcriptional coactivator in RNA polymerase II transcription. This gene contains stretches of trinucleotide repeats and is located in the chromosome 22 region which is deleted in DiGeorge syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2014]</p> <p>Transcript Variant: This variant (3) lacks an alternate in-frame exon and uses an alternate in-frame splice site in the 3' coding region, compared to variant 1. The encoded protein (isoform c) is shorter than isoform a.</p>