

Product datasheet for **SC337397**

MARK1 (NM_001286128) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	MARK1 (NM_001286128) Human Untagged Clone
Tag:	Tag Free
Symbol:	MARK1
Synonyms:	MARK; Par-1c; Par1c
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >NCBI ORF sequence for NM_001286128, the custom clone sequence may differ by one or more nucleotides

```

ATGTCGGCCCGGACGCCATTGCCGACGGTGAACGACGGGACACGAAAAATCATACATCTGTGGATGGAT
ATACTGAACCACACATCCAGCCTACCAAGTCGAGTAGCAGACAGAACATCCCCGGGTGTAGAACTCCAT
TAGCTCAGCAACAGATGAACAGCCTCACATTGGAAATTACCGTTTACAAAAACAATAGGGAAGGGAAAT
TTTGCCAAAGTCAAATTGGCAAGACACGTTCTAACTGGTAGAGAGGTTGCTGTGAAAATAATAGACAAA
CTCAGCTAAATCTACCAGTCTACAAAAGTTATTTGAGAGAAGTACGAATAATGAAGATACTGAATCATCC
TAATATAGGTGAAGTATTTGATTACTTAGTTGCCCATGGAAGAATGAAAGAGAAAAGGCCCGTGCAAAA
TTTAGGCAGATTGTATCTGCTGTACAGTATTGTCATCAAAGTACATTGTTCCACCGTGATCTTAAGGCTG
AAAACCTTCTCCTTGATGGTGATATGAATATTAATTTGCTGACTTTGGTTTTAGTAATGAATTTACAGT
TGGGAACAAATGGACACATTTTGTGGAAGCCACCCATGCTGCTCCGAGCTTTTCCAAGGAAAGAAG
TATGATGGGCCTGAAGTGGATGTGTGGAGTCTGGGCGTCATTCTCTATACATTAGTCAGTGGCTCCTTGC
CTTTTCGATGGCCAGAATTTAAAGGAACTGCGAGAGCGAGTTTTACGAGGGAAGTACCGTATTCCTTCTA
TATGTCCACAGACTGTGAAAATCTTCTGAAGAAATTAATAGTCCCTGAATCCAATAAAGAGAGGCAGCTTG
GAACAAATAATGAAAGATCGATGGATGAATGTTGGTCATGAAGAGGAAGAACTAAAGCCATATACTGAGC
CTGATCCGGATTTCAATGACACAAAAAGAATAGACATTATGGTACCATGGGCTTTGCACGAGATGAAAT
AAATGATGCCTTAATAAATCAGAAGTATGATGAAGTTATGGTACTTATATTCTTCTAGGTAGAAAAACCA
CCTGAATTTGAAGGTGGTGAATCGTTATCCAGTGGAACTTGTGTGAGAGTCCCGGCCAGTAGTGACT
TAAACAACAGCACTCTTCAGTCCCCTGCTCACCTGAAGGTCCAGAGAAGTATCTCAGCAAATCAGAAGCA
GCGGCGTTTCAGTGATCATGCTGGTCCATCCATTCCTCCTGCTGTATCATATACAAAAGACCTCAGGCT
AACAGTGTGAAAAGTGAACAGAAAAGAGGAGTGGGACAAAAGATGTGGCTCGAAAACCTGGCAGCACAAACAG
TTGGATCAAAAAGCGAGATGACTGCAAGCCCTTGTAGGGCCAGAGAGGAAAAAATCTTCAACTATTCC
AAGTAACAATGTGATTCTGGAGGTAGCATGGCAAGAAGGAATACATATGTCTGTGAAAGGACCACAGAT
CGATACGTAGCATTGCAGAAATGGAAGACAGCAGCCTTACGGAGATGTCTGTGAGTAGCATATCTTCTG
CAGGCTCTTCTGTGGCCTCTGCTGTCCCCTCAGCACGACCCCGCCACCAGAAGTCCATGTCCACTTCTGG
TCATCCTATTAAGTCACTGCCAACCATTAAGACGGCTCTGAAGCTTACCGGCTGGTACAACCCAG
AGAGTGCCTGCTGCTTCCCCTCTGCTCACAGTATTAGTACTGCGACTCCAGACCGGACCCGTTTTCCCC
GAGGGAGCTCAAGCCGAAGCACTTCCATGGTGAACAGCTCCGGGAGCGACGCAGCGTTGCTTATAATGG
GCCACCTGCTTACCATCCATGAAACGGGTGCATTTGCACATGCCAGAAGGGGAACGTCAACTGGTATA
ATAAGCAAAATCACATCCAAATTTGTTCCGAGAAGTACATCAGGGGAACCAAAAAGAAAGAGACAAGGAAG
AGGGTAAAGATTCTAAGCCGCTTCTTTGCGGTTACATGGAGTATGAAGACCACTAGTTCAATGGACCC
TAATGACATGATGAGAGAAATCCGAAAAGTGTAGATGCAAAATACTGTGATTATGAGCAAAAAGAGAGA
TTTTTGTCTTCTGTGTCATGGAGACGCTAGACAGGATAGCCTCGTGCAGTGGGAGATGGAAAGTCTGCA
AGTTGCCACGACTGTCACTTAATGGGGTTCGTTCAAGCGAATATCTGGGACATCTATTGCCTTTAAGAA
CATTGCATCAAAAATAGCAAATGAGCTTAAAGCTGTA
    
```

Restriction Sites: SgfI-MluI

ACCN: NM_001286128

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001286128.1 , NP_001273057.1
RefSeq Size:	5225 bp
RefSeq ORF:	2277 bp
Locus ID:	4139
UniProt ID:	Q9P0L2
Cytogenetics:	1q41
Protein Families:	Druggable Genome, Protein Kinase
Gene Summary:	<p>Serine/threonine-protein kinase (PubMed:23666762). Involved in cell polarity and microtubule dynamics regulation. Phosphorylates DCX, MAP2 and MAP4. Phosphorylates the microtubule-associated protein MAPT/TAU (PubMed:23666762). Involved in cell polarity by phosphorylating the microtubule-associated proteins MAP2, MAP4 and MAPT/TAU at KXGS motifs, causing detachment from microtubules, and their disassembly. Involved in the regulation of neuronal migration through its dual activities in regulating cellular polarity and microtubule dynamics, possibly by phosphorylating and regulating DCX. Also acts as a positive regulator of the Wnt signaling pathway, probably by mediating phosphorylation of dishevelled proteins (DVL1, DVL2 and/or DVL3).[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (4) has multiple differences in the coding region but maintains the reading frame, compared to variant 1. This variant encodes isoform 4, which is shorter than isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>