

## Datasheet for NCI-H1299-Cas9 AAVS1 Cell Line

**Catalog number:** SL550001

**Product:** NCI-H1299 cell line stably expressing CRISPR Cas9 nuclease


**Description:** This product is a cell line stably expressing the CRISPR Cas9 nuclease. Cas9 is integrated at the human AAVS1 Safe Harbor locus (also known as PPP1R2C). This cell line also expresses copGFP and the puromycin resistance gene. In combination with separately transfected or transduced single guide RNAs (sgRNAs), this cell line will sustain double-strand DNA breaks (DSBs) at targeted genome sites. This cell line can be used *in vitro* for gene knockout, transgene knockin, mutagenesis, transgene integration, or other genome editing-related applications

**Quantity:** 1 vial of  $2 \times 10^6$  cells; frozen

**Shipping conditions:** Dry ice

**Storage conditions:** Liquid nitrogen vapor phase. Remove the item from the dry ice packaging and check all items for damage and leakage. Place immediately into storage at or below  $-140^{\circ}\text{C}$ , preferably into the liquid nitrogen vapor phase, until use.

**Transgene integration:**



**Source of parental line:**

NCI-1299

Organism: *Homo sapiens*, human

Tissue: lung, derived from metastatic lymph node

Cell type: Epithelial

Pathology: Carcinoma, non-small cell lung cancer



**Safety instructions:** To ensure safety, protective gloves, clothing, and a face mask should be worn when handling frozen vials. Some leakage may occur into the vial during storage. The liquid nitrogen will be converted to gas upon thawing. Due to the nature of nitrogen gas, pressure may build within the vial and possibly result in the vial exploding or losing its cap. This may cause flying debris.

**Thawing procedure:** The vial of cells should be thawed in a 37 °C water bath with gentle agitation. For optimal performance, the vial should be thawed in under two minutes. Ensure that the cap of the vial did not loosen upon thawing, and re-tighten as needed. Spray the vial with 70% EtOH and wipe off. Repeat. Using aseptic technique, add the contents of the vial to 9 ml of complete growth medium (without selection). Centrifuge for 5 min. at 125 x g. Aspirate the medium, being careful not to disturb the pellet. Resuspend in 10 mL of complete growth medium, and place into a culture vessel of your choice. Only add selection to the medium after 24 hours in culture.

**Culture conditions:**

**Complete Growth Medium**

The base medium for this cell line is RPMI1640. For optimal growth and maintenance of selection, add the following components to the base medium: fetal bovine serum to a final concentration of 10%.

**Selection**

Puromycin to a final concentration of 1 µg/mL

**Culture temperature:**

37 °C with 5% CO<sub>2</sub>

**Subculture:**



Rinse the cells with PBS without cations, digest cells with 0.25% (w/v) Trypsin-EDTA (0.53 mM) solution and split at 1:3 to 1:10 ratio.

**Cryopreservation:** Freeze slowly in complete growth medium supplemented with 5% (v/v) DMSO.

**Citation of product:** If use of this item results in a publication, please use this information: CRISPR Cas9 stable NCI-H1299 cell line (SL550001, OriGene Technologies, Inc., Rockville, MD).


**Quality control:** >95% viability before freezing. All cells were tested and found to be free of mycoplasma, bacterial, viruses, and other toxins.

**1. Cas9 gene integration at AAVS1 site in H1299/Cas9 cell line by Junctional PCR from genomic DNA**




**2. H1299/Cas9 Activity by T7 Endonuclease I assay (T7E1)**

sgRNA targeting to EMX1 or HUWE gene were transiently transfected into H1299/CAS9 AAVS1 cell line. EMX1 or HUWE gene was cut by CAS9 expressed inside the cells and repaired through NHEJ with mutation. The mutations will be recognized and cut by T7 Endonuclease I.



For EMX1 sgRNA transient transfected cells, a 684 bp EMX1 gene fragment from PCR was tested by T7E1 Assay. The T7E1 cleavage will results in two additional bands:  
315 bp and 369 bp.



For HUWE sgRNA transient transfected cells, a 525 bp HUWE gene fragment from PCR was tested by T7E1 Assay. The T7E1 cleavage will results in two additional bands:  
192 bp and 333 bp.



### Limited Use License

A limited use license is granted to the Buyer of the Product. The Product shall be used by the Buyer for internal research purposes only and shall not be transferred (in any way) to any third party. The Product is expressly not designed, intended, or warranted for use in humans or for therapeutic or diagnostic use. The Product must not be resold, repackaged or modified for resale, or used to manufacture commercial products without prior written consent from OriGene. This Product should be used in accordance with NIH guidelines developed for recombinant DNA and genetic research. Use of any part of the Product constitutes acceptance of the above terms.