

## Product datasheet for SM1057APC

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **CD3E Mouse Monoclonal Antibody [Clone ID: UCHT1]**

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: UCHT1

Applications: FC

**Recommended Dilution:** Flow Cytometry: Use 10 μl of neat antibody to label 16 cells or 100 μl whole blood

Reactivity: Human, Primate

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human infant thymocytes and lymphocytes from a patient with Sezary Syndrome.

Spleen cells from immunised BALB/c mice were fused with cells of the P3/NS1/1-Ag4-1 mouse

myeloma cell line.

**Specificity:** This antibody recognises a surface protein of peripheral blood T cells of molecular weight

19kD. As a pan T cell marker this antibody is present on more than 95% of circulating human

peripheral T cells.

Formulation: PBS, pH 7.2

Label: APC

State: Liquid purified IgG fraction Stabilizer: 1% BSA, 5% Sucrose Preservative: 0.09% Sodium Azide

Label: Allophycocyanin

**Reconstitution Method:** Restore with distilled water

**Purification:** Affinity Chromatography on Protein G

Conjugation: APC

**Storage:** Prior to and following reconstitution store the antibody undiluted at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

**Stability:** Shelf life: one year from despatch.



#### CD3E Mouse Monoclonal Antibody [Clone ID: UCHT1] - SM1057APC

Database Link: Entrez Gene 916 Human

P07766


**Background:** T Cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of the CD3

subunits CD3 gamma, CD3 delta, CD3 epsilon and CD3 zeta. These CD3 subunits are structurally related members of the immunoglobulins super family encoded by closely linked genes on human chromosome 11. The CD3 components have long cytoplasmic tails that associate with cytoplasmic signal transduction molecules. This association is mediated at least in part by a double tyrosine-based motif present in a single copy in the CD3 subunits. CD3

may play a role in TCR-induced growth arrest, cell survival and proliferation.

Synonyms: T3/Leu-4

# **Product images:**



Staining of human peripheral blood lymphocytes with Mouse ANTI Human CD3-APC