

Product datasheet for SM1057R

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD3E Mouse Monoclonal Antibody [Clone ID: UCHT1]

Product data:

Product Type: Primary Antibodies

Clone Name: UCHT1

Applications: FC

Recommended Dilution: Flow cytometry.

Reactivity: Human, Primate

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human infant thymocytes and lymphocytes from a patient with Sezary Syndrome.

Specificity: This antibody recognises CD3, a 19 kD cell surface protein which is expressed on more than

95% of circulating peripheral blood T-cells.

Formulation: PBS, pH 7.2 containing 0.09% Sodium Azide as preservative and 1% BSA as stabilizer.

Label: PE

State: Lyophilized purified IgG fraction.

Label: R. Phycoerythrin (RPE)

Reconstitution Method: Restore with distilled water.

Concentration: lot specific

Purification: Affinity chromatography on Protein G

Conjugation: PE

Storage: Store the antibody undiluted at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Stability: Shelf life: one year from despatch.

Gene Name: CD3e molecule

Database Link: Entrez Gene 916 Human

P07766






Background:

T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of the CD3 subunits: CD3 gamma, CD3 delta, CD3 epsilon and CD3 zeta. These CD3 subunits are structurally related members of the immunoglobulins super family encoded by closely linked genes on human chromosome 11. The CD3 components have long cytoplasmic tails that associate with cytoplasmic signal transduction molecules. This association is mediated at least in part by a double tyrosine based motif present in a single copy in the CD3 subunits. CD3 may play a role in TCR induced growth arrest, cell survival and proliferation. The CD3 antigen is present on 68-82% of normal peripheral blood lymphocytes, 65-85% of thymocytes and Purkinje cells in the cerebellum. It is never expressed on B or NK cells. Decreased percentages of T lymphocytes may be observed in some autoimmune diseases.

Synonyms: T3/Leu-4

Product images:



Staining of human peripheral blood lymphocytes with Mouse Anti Human CD3-RPE.