

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for SM1057RT

## CD3E Mouse Monoclonal Antibody [Clone ID: UCHT1]

### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	UCHT1
Applications:	FC
Recommended Dilution:	Flow cytometry.
Reactivity:	Human, Monkey
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Human infant thymocytes and lymphocytes from a patient with Sezary Syndrome.
Specificity:	This antibody recognises CD3, a 19 kD cell surface protein which is expressed on more than 95% of circulating peripheral blood T-cells.
Formulation:	PBS, pH 7.2 containing 0.09% Sodium Azide as preservative and 1% BSA as stabilizer. Label: PE State: Lyophilized purified IgG fraction. Label: R. Phycoerythrin (RPE)
Reconstitution Method:	Restore with distilled water.
Concentration:	lot specific
Purification:	Affinity chromatography on Protein G
Conjugation:	PE
Storage:	Store the antibody undiluted at 2-8°C. <b>DO NOT FREEZE!</b> This product is photosensitive and should be protected from light.
Stability:	Shelf life: one year from despatch.
Gene Name:	CD3e molecule
Database Link:	<u>Entrez Gene 916 Human</u> <u>P07766</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** CD3E Mouse Monoclonal Antibody [Clone ID: UCHT1] – SM1057RT

Background:T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of the CD3<br/>subunits: CD3 gamma, CD3 delta, CD3 epsilon and CD3 zeta. These CD3 subunits are<br/>structurally related members of the immunoglobulins super family encoded by closely linked<br/>genes on human chromosome 11. The CD3 components have long cytoplasmic tails that<br/>associate with cytoplasmic signal transduction molecules. This association is mediated at least<br/>in part by a double tyrosine based motif present in a single copy in the CD3 subunits. CD3<br/>may play a role in TCR induced growth arrest, cell survival and proliferation. The CD3 antigen<br/>is present on 68-82% of normal peripheral blood lymphocytes, 65-85% of thymocytes and<br/>Purkinje cells in the cerebellum. It is never expressed on B or NK cells. Decreased<br/>percentages of T lymphocytes may be observed in some autoimmune diseases.

Synonyms: T3/Leu-4

#### **Product images:**



Staining of human peripheral blood lymphocytes with Mouse Anti Human CD3-RPE.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US