

## **Product datasheet for SM1073P**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CD16 (FCGR3A) Mouse Monoclonal Antibody [Clone ID: LNK16]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: LNK16
Applications: FC, IP

**Recommended Dilution:** Flow Cytometry: Use 10 μl of 1/10-1/20 diluted antibody to label 10e6 cells or 100 μl human

whole blood.

Immunoprecipitation.

Does not work in Immunohistochemistry on Frozen and Paraffin Sections.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Normal Human peripheral blood granulocytes.

**Specificity:** This antibody recognizes CD16.

This antibody has been reported to block the binding of IgG to the CD16 molecule. (4)

Formulation: PBS, pH 7.4

State: Purified

State: Liquid purified IgG fraction Preservative: 0.09% Sodium Azide

**Concentration:** lot specific

**Purification:** Ion Exchange Chromatography

**Conjugation:** Unconjugated

**Storage:** Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

**Gene Name:** Fc fragment of lgG receptor Illa

Database Link: Entrez Gene 2214 Human

P08637





## CD16 (FCGR3A) Mouse Monoclonal Antibody [Clone ID: LNK16] - SM1073P

**Background:** CD16 is a 50-65kD cell surface molecule which exists in two forms, a transmembranous form

expressed by NK cells and some T cells, and a phosphatidylinositol linked form expressed by granulocytes. CD16 is a low affinity receptor for IgG (FcR III), and is an important receptor

mediating ADCC by NK cells.

Synonyms: FCGR3A, CD16A, FCG3, FCGR3, IGFR3, Fc-gamma RIII-alpha, Fc-gamma RIII, Fc-gamma RIIIa,

FcRIII, FcRIIIa, FcR-10, IgG Fc receptor III-2