

Product datasheet for SM1097AS

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD35 (CR1) Mouse Monoclonal Antibody [Clone ID: E11]

Product data:

Product Type: Primary Antibodies

Clone Name: E11

Applications: FC, IHC, WB

Recommended Dilution: Flow Cytometry: 1/50 - 1/100.

Immunohistochemistry on Frozen Sections.

Immunohistochemistry on Paraffin Sections: 10 µg/ml.

Western Blot.

Reactivity: Human, Monkey, Baboon

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human acute monocytic leukaemia cells.

Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma

cell line.

Specificity: This antibody is a single chain cell surface glycoprotein which exists in four allotypic forms

(A,B,C,D) of 190kD, 220kD, 160kD and 250kD respectively. The molecule is expressed by granulocytes, monocytes, B cells and some T cells. Functionally CD35 acts as a receptor for

the C3b component of complement.

Formulation: PBS, pH 7.4

State: Azide Free

State: Liquid purified IgG fraction from Tissue Culture Supernatant

Stabilizer: None Preservative: None

Concentration: lot specific

Purification: Affinity Chromatography on Protein A

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.





CD35 (CR1) Mouse Monoclonal Antibody [Clone ID: E11] - SM1097AS

Gene Name: complement component 3b/4b receptor 1 (Knops blood group)

Database Link: Entrez Gene 718120 MonkeyEntrez Gene 1378 Human

P17927

Background: CD35 is a membrane glycoprotein found on peripheral blood cells, glomerular podocytes,

and follicular dendritic cells. The protein is a receptor for complement components C3b and C4b and regulates the activity of the complement cascade. Variation in this protein is the basis of the Knops blood group system. The two most common alleles, F and S, differ by 8 exons and are thought to be the result of an unequal crossover event. A secreted form of the

protein present in plasma has been described, but its full length nature has not been


determined.

Synonyms: Complement receptor type 1, CR1, C3b/C4b receptor

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Complement and coagulation cascades, Hematopoietic cell lineage

Product images:



Formalin-Fixed, Paraffin-Embedded Human Kidney stained with Azide Free CR1 Antibody (Clone E11) 10 ug/ml after heat-induced antigen retrieval.