

Product datasheet for SM1125PT

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD49b (ITGA2) Mouse Monoclonal Antibody [Clone ID: AK7]

Product data:

Product Type: Primary Antibodies

Clone Name: AK7

Applications: FC, IHC

Recommended Dilution: Flow Cytometry: Use 10 μl of 1/50-1/100 diluted antibody to label 10e6 cells in 100 μl.

Immunohistochemistry on Frozen Sections: 1/500-1/1000.

Recommended Positive Control Tissue: Tonsil.

Reactivity: Human, Monkey

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Spleen cells from immunised mice were fused with cells of the mouse NS1 myeloma cell line.

Specificity: This antibody recognises the integrin alpha 2 subunit, which is 160kD cell surface antigen

expressed by NK cells and a small subset of T-cells. This antibody inhibits cell attachment to collagen.

Formulation: PBS containing 0.09% Sodium Azide as preservative.

State: Purified

State: Liquid purified IgG fraction.

Concentration: lot specific

Purification: Affinity Chromatography on Protein G.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: integrin subunit alpha 2

Database Link: Entrez Gene 3673 Human

P17301






Background:

Integrins are heterodimeric cell surface receptors composed of alpha and beta subunits, which mediate cell-cell and cell-extracellular matrix attachments. They are responsible for adhesion of platelets and other cells to collagens, modulation of collagen and collagenase gene expression, force generation and organization of newly synthesized extracellular matrix. Aberrant integrin expression has been found in many epithelial tumours. Changes in integrin expression have been shown to be important for the growth and early metastatic capacity of melanoma cells.

Synonyms:

Integrin alpha-2, ITGA-2, VLA-2 alpha, VLA2, GPIa, Collagen Receptor

Product images:



Staining of human peripheral blood monocytes with Mouse Anti Human CD49b.