

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for SM1140FT

## Integrin alpha 6 (ITGA6) Mouse Monoclonal Antibody [Clone ID: 450-30A]

### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	450-30A
Applications:	FC
Recommended Dilution:	<b>Flow Cytometry:</b> Use 10 $\mu$ l of neat antibody to label 10e6 cells in 100 $\mu$ l.
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Alpha 6 beta 4 Integrin purified from A431 cells
Specificity:	This antibody recognizes the human VLA-6 cell surface antigen, also known as the a6 Integrin and as CD49f. CD49f is expressed by platelets, weakly by monocytes and by a subset of lymphocytes.
Formulation:	PBS, pH 7.4 Label: FITC State: Liquid purified IgG fraction Stabilizer: 1% BSA Preservative: 0.09% Sodium Azide Label: Fluorescein Isothiocyanate Isomer 1
Concentration:	lot specific
Purification:	Affinity chromatography on Protein G
Conjugation:	FITC
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	integrin subunit alpha 6
Database Link:	Entrez Gene 3655 Human P23229




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### CRIGENE Integrin alpha 6 (ITGA6) Mouse Monoclonal Antibody [Clone ID: 450-30A] – SM1140FT

Background:Integrins are important extracellular matrix (ECM) receptor proteins located on cell surfaces.<br/>They are hetrodimers composed of an alpha and a beta transmembrane glycoprotein<br/>subunit. Around twenty two different integrins (different alpha/ beta subunit combinations)<br/>are found in nature. Integrins are generally present in high concentrations at the cell surface,<br/>but, unlike most other cell surface receptors, they bind ligands with very low affinity. Due to<br/>their weak individual binding, integrins need to cluster and bind in groups in order to<br/>effectively bind the ECM. Integrins bind many different ligands including laminin. Each<br/>integrin is made up of a large N terminal extracellular domain that binds the ECM ligand and<br/>a small C terminal cytoplasmic domain that mediates interaction with the actin cytoskeleton<br/>and signaling function. Integrin alpha 6 complexes are receptors for laminins, which are<br/>components of basement membranes. Integrin alpha 6 complexes may play an important<br/>role in embryogenesis.

Synonyms: Integrin alpha-6, VLA-6, VLA6

#### **Product images:**



Staining of human peripheral blood lymphocytes with Mouse Anti Human CD49f.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US