

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for SM1149R

CD62L (SELL) Mouse Monoclonal Antibody [Clone ID: FMC46]

Product data:

Product Type:	Primary Antibodies
Clone Name:	FMC46
Applications:	FC
Recommended Dilution:	Flow Cytometry: Use 10 µl of neat antibody to label 1౷ cells or 100 µl whole blood.
Reactivity:	Bovine, Human, Monkey, Canine
Host:	Mouse
lsotype:	lgG2b
Clonality:	Monoclonal
Immunogen:	PHA stimulated lymphoblasts. Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
Specificity:	This antibody recognizes CD62L, also known a L-selectin.
Formulation:	PBS Label: PE State: Lyophilized purified IgG Stabilizer: 5% Sucrose, 1% BSA Preservative: 0.09% Sodium Azide Label: R. Phycoerythrin (RPE)
Reconstitution Method:	Restore with 1 ml distilled water.
Purification:	Affinity chromatography on Protein G
Conjugation:	PE
Storage:	Store undiluted at 2-8°C. DO NOT FREEZE! This product is photosensitive and should be protected from light.
Stability:	Shelf life: one year from despatch.
Gene Name:	selectin L
Database Link:	<u>Entrez Gene 6402 Human</u> <u>P14151</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	CD62L (SELL) Mouse Monoclonal Antibody [Clone ID: FMC46] – SM1149R
Background:	CD62L is a 74-95kDa member of the selectin family of adhesion receptors, which acts as a ligand for both CD62-P (P-selectin) and CD62E (E-selectin). Human CD62L is constitutively expressed on most leucocytes including monocytes, granulocytes, lymphocytes, NK cells, bone marrow myeloid progenitor cells and on a subset of thymocytes. CD62L plays an important role in leucocyte tethering and rolling on the endothelial cell surface and for the homing of naive lymphocytes to lymph nodes and Peyers patches via HEV. Neutrophils require a constant supply of this molecule on the cell surface for migration into peripheral tissues and adhesion to activated endothelium at sites of inflammation, where CD62L is rapidly shed as soluble L-selectin, but surface expression still remains. The expression of CD62L is down regulated on lymphocytes and neutrophils by PMA stimulation.
Synonyms:	SELL, LNHR, LYAM1, Leu-8, TQ1, gp90-MEL, LECAM1, LAM-1

%// _

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US