

Product datasheet for SM1150P

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD62P (SELP) Mouse Monoclonal Antibody [Clone ID: AK-6]

Product data:

Product Type: Primary Antibodies

Clone Name: AK-6

Applications: FC, IHC, WB

Recommended Dilution: Western blotting.

Flow Cytometry: Use 10 μl of 1/10-1/100 diluted antibody to label 16 cells in 100 μl.

Immunohistochemistry on Frozen Sections: 1/1000-1/5000.

Immunohistochemistry on Paraffin Embedded Sections: 1/500-1/2000. This product does not require antigen retrieval using heat treatment prior to staining of paraffin sections but

staining may be enhanced using Sodium Citrate buffer pH 6.0.

Recommended Positive Control Tissue: Tonsil.

Reactivity: Human, Monkey

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human platelet membrane glycoproteins.

Specificity: This antibody recognizes CD62P.

It reacts with a 140 kDa antigen expressed on activated platelets and also on endothelial cells

both of blood and lymph vessels.

Formulation: PBS, pH 7.4

State: Purified

State: Liquid purified IgG fraction from Tissue Culture Supernatant

Preservative: 0.09% Sodium Azide

Concentration: lot specific

Purification: Affinity Chromatography on Protein G

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.





Gene Name: selectin P

Database Link: Entrez Gene 6403 Human


P16109

Background: CD62P (P-Selectin) is an adhesion glycoprotein that is expressed on platelets and endothelial


cells upon their activation. Interaction between CD62P and its mucin-like ligand PSGL-1 (P-selectin glycoprotein ligand-1) expressed on the microvilli of most leukocytes supports leukocyte rolling along postkapillary venules at the earliest time of inflammation. Both CD62P and PSGL-1 are extended glycoproteins that form homodimers. CD62P dimerization is probably mediated through interactions of the transmembrane domains and stabilizes leukocyte tethering and rolling, probably by increasing rebinding within a bond cluster.

Synonyms: SELP, GMRP, GRMP, PADGEM, GMP-140, LECAM3

Product images:



Staining of Human tonsil showing capillary endothelium. Formalin fixed, Paraffin processed tissue with Mouse ant-Human CD62P Antibody.



Staining of Human peripheral blood platelets with anti-Human CD62P Antibody