

Product datasheet for SM1166PS

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD71 (TFRC) Mouse Monoclonal Antibody [Clone ID: DF1513]

Product data:

Product Type: Primary Antibodies

Clone Name: DF1513

Applications: FC, IF, IHC, WB

Recommended Dilution: Flow Cytometry: use 10 μl of 1/10-1/25 diluted antibody to label 1x10e6 cells in 100 μl.

Immunofluorescence.

Immunohistochemistry on Frozen Tissue Sections only: 10-20 µg/ml with an avidin-biotin

system.

Western blot: 1-5 µg/ml.

Reactivity: Human, Monkey

Host: Mouse Isotype: IgG1

Clonality: Monoclonal Immunogen: KGI cell line.

Specificity: This antibody recognizes the Human CD71 cell surface antigen, a 190 kD homodimeric

glycoprotein expressed by proliferating cells.

Formulation: PBS containing 0.09% Sodium Azide as preservative

State: Aff - Purified

State: Liquid purified IgG fraction

Concentration: lot specific

Purification: Affinity Chromatography on Protein G

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: transferrin receptor

Database Link: Entrez Gene 7037 Human

P02786





CD71 (TFRC) Mouse Monoclonal Antibody [Clone ID: DF1513] - SM1166PS

Background:

The transferrin receptor has been structurally characterized as a sulfide bound dimer of identical glycoprotein subunits of 95 kDa. The transferrin receptor is not present on resting blood lymphocytes. On PBL, the receptor appears after activation. The expression of transferrin receptor is coordinately regulated with cell growth. Present on T and B cell lines. The soluble (or serum) transferrin receptor (sTfR) is a circulating truncated form of the membrane receptor protein; it is an 85 kDa glycoprotein forming in serum a 320 kDa complex with diferric transferrin. The most important clinical use of the sTfR determination is in the differential diagnosis between iron deficiency anaemia and the anaemia of chronic disease.

Synonyms:

TfR1, p90, Transferrin receptor protein 1