

Product datasheet for SM1166RT

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD71 (TFRC) Mouse Monoclonal Antibody [Clone ID: DF1513]

Product data:

Product Type: Primary Antibodies

Clone Name: DF1513

Applications: FC

Recommended Dilution: Flow Cytometry: Use 10 µl of neat antibody to label 1x10e6 cells.

Reactivity: Human, Monkey

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: KGI cell line. Spleen cells from immunised BALB/c mice were fused with cells of the mouse

NS1 myeloma cell line.

Specificity: This antibody recognises the human CD71 cell surface antigen, a 190 kD homodimeric

glycoprotein expressed by proliferating cells. CD71 acts as receptor for transferrin.

Formulation: PBS, pH 7.4 containing 0.09% Sodium Azide as preservative and 0.1% BSA as stabilizer.

Label: PE

State: Lyphilized purified IgG fraction.

Label: R-Phycoerythrin

Reconstitution Method: Restore with 0.25 ml distilled water.

Concentration: lot specific

Purification: Affinity Chromatography on Protein G.

Conjugation: PE

Storage: Prior to and following reconstitution store the antibody undiluted at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Stability: Shelf life: one year from despatch.

Gene Name: transferrin receptor

Database Link: Entrez Gene 7037 Human

P02786






Background:

The transferrin receptor has been structurally characterized as a sulfide bound dimer of identical glycoprotein subunits of 95 kDa. The transferrin receptor is not present on resting blood lymphocytes. On PBL, the receptor appears after activation. The expression of transferrin receptor is coordinately regulated with cell growth. Present on T and B cell lines. The soluble (or serum) transferrin receptor (sTfR) is a circulating truncated form of the membrane receptor protein; it is an 85 kDa glycoprotein forming in serum a 320 kDa complex with diferric transferrin. The most important clinical use of the sTfR determination is in the differential diagnosis between iron deficiency anaemia and the anaemia of chronic disease.

Synonyms:

TfR1, p90, Transferrin receptor protein 1

Product images:



Staining of Jurkat cells with Mouse Anti Human CD71-RPE.