

## Product datasheet for SM1602PS

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### CD147 (BSG) Mouse Monoclonal Antibody [Clone ID: MEM-M6/1]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: MEM-M6/1

**Applications:** FC, IHC, IP, WB

Recommended Dilution: Flow Cytometry: 10 µg/ml.

Immunoprecipitation.

**Western blot** (under non-reducing conditions): 1 μg/ml. **Immunohistochemistry on Paraffin Sections:** 10 μg/ml.

Positive tissue: testis.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Protein A-CR purified soluble recombinant form of CD147, CD147Rg, which consists of the

cDNA coding for the hinge region, CH2-and CH3 domain of Human IgG1 (CD147Rg is secreted

by transfectants as a dimer)

**Specificity:** The antibody recognizes an epitope in the N-terminal Ig domain (D1) of CD147 (Neurothelin),

a 50-60 kDa type I transmembrane glycoprotein primarily expressed on all leukocytes, red blood cells, platelets and endothelial cells; it is not expressed by resting lymphocytes. The antibody MEM-M6/1 is a high-affinity antibody capable of binding to unstimulated

peripheral blood T cells.

**Formulation:** PBS, pH~7.4

State: Purified

State: Liquid purified Ig fraction (> 95% by SDS-PAGE)

Preservative: 15 mM Sodium Azide

**Concentration:** lot specific

**Purification:** Affinity Chromatography on Protein A

**Conjugation:** Unconjugated

**Storage:** Store undiluted at 2-8°C.

DO NOT FREEZE!





#### CD147 (BSG) Mouse Monoclonal Antibody [Clone ID: MEM-M6/1] - SM1602PS

**Stability:** Shelf life: one year from despatch.

Database Link: Entrez Gene 682 Human

P35613

Background: CD147 (basigin, neurothelin, OX-47, 5A11, CE9, M6) also known as EMMPRIN (extracellular

matrix metalloproteinase inducer) or TCSF (tumour cell-derived collagenase-stimulatory factor) is an ubiquitously expressed cell surface protein with multiple glycosylated forms. The highest level of CD147 expression is on metabolically active cells, such as lymphoblasts, inflammatory cells, brown adipocytes and malignant tumour cells. CD147 has multiple functions, including facilitating of cell surface expression of monocarboxylate transporter proteins and extracellular matrix metalloproteinases, regulation of integrin functions, it plays

roles in cell development and activation, fetal development or retinal function.

Synonyms: Basigin, BSG, EMMPRIN, TCSF