

Product datasheet for SM1795B

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD45 (PTPRC) Mouse Monoclonal Antibody [Clone ID: F-10-89-4]

Product data:

Product Type: Primary Antibodies

Clone Name: F-10-89-4

Applications: FC

Recommended Dilution: Flow cytometry.

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human T lymphocytes. Spleen cells from immunised BALB/c mice were fused with cells of the

mouse NS-1 myeloma cell line.

Specificity: This antibody recognises the CD45 cell surface antigen, also known as the leucocyte common

antigen (LCA). Antibodies recognising a common epitope on all of these isoforms are termed CD45 whilst those recognising only individual isoforms are termed CD45RA or CD45RO etc. Clone F10-89-4 reacts with all forms of CD45 expressed by all haematopoietic cells, except erythrocytes, having a higher level of expression on lymphocytes than on granulocytes.

Formulation: PBS, pH7.4 containing 0.09% Sodium Azide and 1% Bovine Serum Albumin

Label: Biotin

State: Liquid purified IgG

Concentration: lot specific

Purification: Affinity chromatography on Protein G

Conjugation: Biotin

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: protein tyrosine phosphatase, receptor type C

Database Link: Entrez Gene 5788 Human

P08575






Background:

CD45 is a family of single chain transmembraneous glycoproteins consisting of at least four isoforms (220, 205, 190, 180 kDa) which share a common large intracellular domain. Their extracellular domains are heavily glycosylated. The different isoforms are produced by alternative messenger RNA splicing of three exons of a single gene on chromosome 1. CD45 is expressed on cells of the human hematopoietic lineage (including hematopoietic stem cells) with the exception of mature red cells. It is not detected on differentiated cells of other tissues. It is likely that CD45 plays an important role in signal transduction, inhibition or upregulation of various immunological functions. Antibodies recognising a common epitope on all of the isoforms are termed CD45 whilst those recognising only individual isoforms are termed CD45RA or CD45RO etc.

Synonyms:

PTPRC, Leukocyte common antigen, L-CA, T200

Product images:



Staining of human peripheral blood lymphocytes with MOUSE ANTI HUMAN CD45:BIOTIN.