

Product datasheet for **SM1800FT**

Integrin alpha E (ITGAE) Mouse Monoclonal Antibody [Clone ID: LF61]

Product data:

| | |
|-----------------------|---|
| Product Type: | Primary Antibodies |
| Clone Name: | LF61 |
| Applications: | FC |
| Recommended Dilution: | Flow Cytometry: Use 5 µl of neat-1/10 diluted antibody to label 10 ⁶ cells in 100 µl. |
| Reactivity: | Human |
| Host: | Mouse |
| Isotype: | IgG1 |
| Clonality: | Monoclonal |
| Immunogen: | Hairy cell leukaemia cells. Spleen cells from immunised BALB/c mice were fused with cells of the mouse NSO myeloma cell line. |
| Specificity: | This antibody recognizes the Human CD103 cell surface antigen. Other species not tested. |
| Formulation: | PBS Label: FITC State: Liquid purified IgG fraction Stabilizer: 0.5% BSA Preservative: 0.09% Sodium Azide Label: Fluorescein Isothiocyanate Isomer 1 Absorption emission: 490 nm / 525 nm |
| Concentration: | lot specific |
| Purification: | Affinity Chromatography on Protein A |
| Conjugation: | FITC |
| Storage: | Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing. |
| Stability: | Shelf life: one year from despatch. |
| Gene Name: | integrin subunit alpha E |



[View online »](#)

Database Link: [Entrez Gene 3682 Human P38570](#)

Background: CD103/Integrin alpha E is a new member of the integrin series of adhesion molecules containing the beta7 subunit. This antigen defines a developmentally important subset of T cells, namely mucosal T cells including all IEL (intraepithelial lymphocytes) and ~20% of lamina propria T cells. Expression of CD103 is more restricted outside these mucosal organs, appearing at lower levels on T cell subsets of the lymph node, dendritic epidermis and periphery. In non-epithelial CD103+ T cells there is a bias toward expression on CD8+ cells. Integrin alpha E is a heterodimer comprising a 150kDa alpha chain and a 120kDa beta chain. Integrin alpha E is expressed by approximately 1% of peripheral blood lymphocytes, activated T lymphocytes and by hairy cell leukaemia cells. The antigen is also expressed by intraepithelial lymphocytes. It has recently been shown to be identical to the alpha E integrin.

Synonyms: Integrin alpha-E, Integrin alpha-IEL, HML-1 antigen, ITGAE