

## Product datasheet for SM2096HRP

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Rat Lambda light chain Mouse Monoclonal Antibody [Clone ID: MARL-15]

## **Product data:**

**Product Type:** Primary Antibodies

Clone Name: MARL-15

**Applications:** ELISA

Recommended Dilution: ELISA.

**Reactivity:** Rat

**Host:** Mouse

**Isotype:** IgG1

Clonality: Monoclonal

Immunogen: Rat lambda light chain. Spleen cells from immunised BALB/c mice were fused with cells of the

rat PAI-O myeloma cell line.

**Specificity:** This antibody recognises immunoglobulin lambda light chain. This antibody shows no cross-

reactivity to Ig heavy chains or to kappa light chain.

**Formulation:** PBS, pH7.4 containing 0.01% Thiomersal and 50% Glycerol

Label: HRP

State: Liquid purified IgG Label: Horseradish Peroxidase

**Concentration:** lot specific

**Purification:** Affinity chromatography

Conjugation: HRP

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.







Background:

All five immunoglobulin classes share the same basic four polypeptide chain structure of two heavy-chains and two light chains. There are five heavy chain types, and two light-chain types (Kappa and Lambda) both having a molecular weight of 22.5kDa. Any heavy-chain type can associate with either light-chain type, but on any immunoglobulin molecule both light-chains are of the same type. Kappa and Lambda consist of a variable region and a constant region and can easily be differentiated by the antigenic properties of the constant region. The ratio of Kappa to Lambda is 70:30, the vast majority of which is bound to heavy-chain in immunoglobulin. In normal individuals low levels of free light-chain arepresent in serum (kappa, 1.6-15.2 mg/L; Lambda, 0.4-4.2mg/L), with the occurrence of multiple myeloma or other B-cell malignancies these levels can be greatly elevated and can be found at high levels in the urine (Bence-Jones proteins).