

## Product datasheet for SM2207F

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **CD8A Mouse Monoclonal Antibody [Clone ID: CVS8]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: CVS8
Applications: FC

**Recommended Dilution:** Flow Cytometry: Use 10 μl of neat-1/10 diluted CD8 antibody to label 16 cells in 100 μl.

**Reactivity:** Donkey, Equine

Host: Mouse Isotype: IgG1

Clonality: Monoclonal Immunogen: Equine PBMC.

Fusion Partners: Spleen cells from immunised BALB/c mice were fused with cells of the X63-

Ag8.653 myeloma cell line.

**Specificity:** This monoclonal CVS8 antibody recognises the Equine homologue of the Human CD8 cell

surface antigen which is expressed by a subset of T lymphocytes.

A study undertaken using CVS8 to identify CD8 on several wild african equid species indicates

that the CVS8 clone recognises Somali wild ass (Equus asinus)  ${f but\ not}$  Grévy's Zebra (E.

grevyi) or Hartmann's Mountain Zebra (E. zebra) (Ibrahim, S, 2007).

Formulation: PBS

Label: FITC

State: Liquid purified IgG fraction from tissue Culture Supernatant

Stabilizer: 1% BSA

Preservative: 0.09% Sodium Azide

**Concentration:** lot specific

**Purification:** Affinity Chromatography on Protein A

Conjugation: FITC

**Storage:** Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.





Gene Name: CD8a molecule

Database Link: P01732


**Background:** The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that

mediates efficient cell to cell interactions within the immune system. The CD8 antigen, acting as a coreceptor, and the T cell receptor on the T lymphocyte recognize antigen displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The functional coreceptor is either a homodimer composed of two alpha chains, or a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant

homology to immunoglobulin variable light chains.

Synonyms: CD8 alpha chain, CD8A, MAL

# **Product images:**



Staining of Equine peripheral blood lymphocytes with Mouse anti Horse CD8 antibody FITC conjugated (Clone CVS8).