

Product datasheet for SM2207P

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD8A Mouse Monoclonal Antibody [Clone ID: CVS8]

Product data:

Product Type: Primary Antibodies

Clone Name: CVS8

Applications: FC, IHC, IP

Recommended Dilution: Flow Cytometry: Use 10 μl of 1/25-1/200 diluted CD8 antibody to label 16 cells in 100 μl.

Immunoprecipitation.

Immunohistochemistry on Frozen Sections.

Reactivity: Donkey, Equine

Host: Mouse Isotype: IgG1

Clonality: Monoclonal Immunogen: Equine PBMC.

Fusion Partners: Spleen cells from immunised BALB/c mice were fused with cells of the X63-

Ag8.653 myeloma cell line.

Specificity: This monoclonal CVS8 antibody recognises the Equine homologue of the Human CD8 cell

surface antigen which is expressed by a subset of T lymphocytes.

A study undertaken using CVS8 to identify CD8 on several wild african equid species indicates that the CVS8 clone recognises Somali wild ass (Equus asinus) **but not** Grévy's Zebra (E.

grevyi) or Hartmann's Mountain Zebra (E. zebra) (Ibrahim, S, 2007).

Negatives Species: Zebra.

Formulation: PBS

State: Purified

State: Liquid purified IgG fraction from tissue Culture Supernatant

Preservative: 0.09% Sodium Azide

Concentration: lot specific

Purification: Affinity Chromatography on Protein A

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.





Gene Name: CD8a molecule

Database Link: P01732


Background: The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that

mediates efficient cell to cell interactions within the immune system. The CD8 antigen, acting as a coreceptor, and the T cell receptor on the T lymphocyte recognize antigen displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The functional coreceptor is either a homodimer composed of two alpha chains, or a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant

homology to immunoglobulin variable light chains.

Synonyms: CD8 alpha chain, CD8A, MAL

Product images:



Staining of Equine peripheral blood lymphocytes with Mouse anti Horse CD8 antibody (Clone CVS8) followed by HRP conjugated Goat anti Mouse IgG HRP