

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for SM2260LE

KLRC4-KLRK1 Mouse Monoclonal Antibody [Clone ID: 1D11]

Product data:

Product Type:	Primary Antibodies
Clone Name:	1D11
Applications:	FC, FN, IHC, IP
Recommended Dilution:	Flow Cytometry: 1/50 - 1/200; use 10 μl of the suggested working dilution to label 106 cells in 100 μl. Immunoprecipitation. Immunohistochemistry on frozen sections. Functional assays.
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	NKL cells. Spleen cells from immunised RBF/DnJ mice were fused with cells of the p3 mouse myeloma cell line.
Specificity:	This antibody is specific for CD314, also known as natural killer receptor G2 (NKG2D) and as killer cell lectin-like receptor subfamily K, member 1 (KLRK1). Clone 1D11 is reported to inhibit T cell recognition of MICA (1,6).
Formulation:	PBS, pH7.4 State: Low Endotoxin State: Liquid purified IgG
Concentration:	lot specific
Purification:	Affinity chromatography on Protein G
Conjugation:	Unconjugated
Storage:	Store the antibody at -20°C. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	KLRC4-KLRK1 readthrough
Database Link:	<u>Entrez Gene 100528032 Human</u> <u>P26718</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	KLRC4-KLRK1 Mouse Monoclonal Antibody [Clone ID: 1D11] – SM2260LE
Background:	CD314 is a C-type lectin-like activating receptor which is expressed on most natural killer (NK) cells, CD8 T cells and gamma delta T cells. CD314 forms homodimers that signal through an associated DAP10 adaptor protein. Ligands of CD314 include MICA, MICB and UL16 binding protein (ULBP), which are inducibly expressed. Ligand binding to CD314 results in NK cell activation and potent costimulation of effector T cells.
Synonyms:	D12S2489E, NKG2D, NK cell receptor D
Note:	Endotoxin Level: less than 0.01EU/mg

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US