

# Product datasheet for SM259F

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Cd8a Mouse Monoclonal Antibody [Clone ID: OX-8]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OX-8
Applications: FC

Recommended Dilution: Flow cytometry: Use 10µl of Neat-1/10 diluted antibody to label 10e6 cells in 100 µl.

This clone has been described reacting with paraffin-embedded material following PLP

Fixation (periodatelysine paraformaldehyde).

Reactivity: Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Rat thymocyte membrane glycoproteins.

Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma

cell line.

**Specificity:** This antibody recognises the CD8 alpha cell surface antigen, expressed by a subset of T

lymphocytes, most thymocytes and the majority of NK cells. Clone MRC OX-8 is suitable for use in in vitro blocking studies.

**Formulation:** PBS, pH 7.4 containing 0.09% Sodium Azide as preservative and 1% BSA as stabilizer.

Label: FITC

State: Liquid purified IgG fraction.

Label: Fluorescein Isothiocyanate Isomer 1

**Concentration:** lot specific

**Purification:** Ion Exchange Chromatography

Conjugation: FITC

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Gene Name: CD8a molecule





### Cd8a Mouse Monoclonal Antibody [Clone ID: OX-8] - SM259F

Database Link: Entrez Gene 24930 Rat

P07725

**Background:** The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that

mediates efficient cell to cell interactions within the immune system. The CD8 antigen, acting as a coreceptor, and the T cell receptor on the T lymphocyte recognize antigen displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The functional coreceptor is either a homodimer composed of two alpha chains, or a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant

homology to immunoglobulin variable light chains.

Synonyms: CD8 alpha chain, CD8A, MAL