

Product datasheet for **SM268FX**

Cd28 Mouse Monoclonal Antibody [Clone ID: JJ319]

Product data:

Product Type:	Primary Antibodies
Clone Name:	JJ319
Applications:	FC
Recommended Dilution:	Flow Cytometry.
Reactivity:	Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Rat CD28 transfected A20J cells Donor: BALB/c spleen Fusion Partner: X63-Ag 8.653
Specificity:	Anti-rat CD28 monoclonal antibody recognizes a 90kDa homeodimeric cell surface glycoprotein. CD28 has been found to be a potent costimulatory receptor on T cells. It is expressed on all peripheral rat ab and most gd T cells, as well as on approximately half of all NK cells. This clone can costimulate T cell proliferation and IL-2 secretion by resting rat T cells.
Formulation:	PBS containing 0.02% sodium azide (NaN ₃) as preservative and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml. Label: FITC State: Liquid purified Ig fraction Label: Fluorescein isothiocyanate isomer 1
Concentration:	lot specific
Purification:	Affinity chromatography on Protein G
Conjugation:	FITC
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	Cd28 molecule


[View online »](#)

Database Link: [Entrez Gene 25660 Rat P31042](#)

Synonyms: TP44

Note: Protocol: **FLOW CYTOMETRY ANALYSIS:**

Method:

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte®-Rat cell separation medium.
2. Wash 2 times.
3. Resuspend the cells to a concentration of 2×10^7 cells/ml in media A. Add 50 μ l of this suspension to each tube (each tube will then contain 1×10^6 cells, representing 1 test).
4. To each tube, add 1.0 μ g* of this antibody per 10^6 cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C. (It is recommended that the tubes are protected from light, since most fluorochromes are light sensitive.)
7. Wash 2 times at 4°C.
8. Resuspend the cell pellet in 50 μ l ice cold media B.
9. Transfer to suitable tubes for flow cytometric analysis containing 15 μ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 μ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100 μ l of 2M sodium azide in 100 mls).

Results:

Tissue Distribution by Flow Cytometry Analysis:

Rat Strain: Fischer

Cell Concentration: 1×10^6 cells per test

Antibody Concentration Used: 1.0 μ g/ 10^6 cells

Isotypic Control: FITC Mouse IgG1

Percentage of cells stained above control:

Thymus 36.3%

Splenic T Cells* 79.7 %

*(T cells isolated with Rat T Cell Recovery Column Kit)

Product images:

