

Product datasheet for SM271PS

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD43 / Leukosialin Mouse Monoclonal Antibody [Clone ID: W3/13]

Product data:

Product Type: Primary Antibodies

Clone Name: W3/13
Applications: FC, IHC

Recommended Dilution: Flow Cytometry: Use 10 μl of 1/50-1/100 diluted antibody to label 10e6 cells in 100 μl.

This product is routinely tested in flow cytometry on rat splenocytes).

Immunohistochemistry on Frozen and Paraffin Sections (requires antigen retrieval using

heat treatment prior to staining).

Reactivity: Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Rat thymocyte membrane glycoproteins

Specificity: SM271PS recognises the Rat CD43 cell surface antigen, also known as leukosialin and

sialophorin. CD43 is a 95kD heavily glycosylated glycoprotein expressed by all leucocytes with

the exception of B lymphocytes.

Formulation: Phosphate buffered saline pH 7.2 with 0.09% Sodium Azide as preservative.

State: Purified

State: Liquid purified IgG fraction

Concentration: lot specific

Purification: Protein G Chromatography

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Should this product contain a precipitate we recommend microcentrifugation before use.

Stability: Shelf life: one year from despatch.

Database Link: P13838





CD43 / Leukosialin Mouse Monoclonal Antibody [Clone ID: W3/13] - SM271PS

Background:

CD43 is one of the major glycoproteins of thymocytes and T lymphocytes, it plays a role in the physicochemical properties of the T cell surface and in lectin binding. CD43 presents carbohydrate ligands to selectins. It has an extended rodlike structure that could protrude above the glycocalyx of the cell and allow multiple glycan chains to be accessible for binding. The antigen is a counter receptor for SN/Siglec1. During T cell activation CD43 is actively removed from the T cell antigen presenting cell contact site suggesting a negative regulatory role in adaptive immune response.

CD43 is expressed by the majority of leucocytes, with the exception of most B cells. However CD43 expression is seen in some cases of B cell lymphocytic lymphoma and centrocytic lymphoma.

Synonyms:

Leukocyte sialoglycoprotein, Sialophorin, Galactoglycoprotein, SPN