

Product datasheet for SM272LE

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Pecam1 Mouse Monoclonal Antibody [Clone ID: TLD-3A12]

Product data:

Product Type: Primary Antibodies

Clone Name: TLD-3A12

Applications: ELISA, FC, FN, IHC, WB

Recommended Dilution: ELISA.

Western Blot.

Flow Cytometry: Use 10 µl of 1/50-1/100 diluted antibody to label 10e6 cells in 100 µl.

Immunohistochemistry on Frozen Sections.

This clone has been reported as suitable for use in IHC onformalin-fixed paraffin-

embedded sections pre-treated with 0.2M boric acid (Wilson, E. et al. (2007)

J.Immunol.Methods. 322: 137-42. Useful in **functional studies**.

Reactivity: Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Activated, Lewis Rat derived microglial cells.

Specificity: This antibody recognises PECAM-1 (CD31).

Clone TLD-3A12 has been shown to partially block the proliferative response of antigen-

specific CD4+ T cells to antigen-presenting cells and relevant antigen.

Formulation: PBS without preservatives or stabilizers

State: Low Endotoxin

State: Liquid purified IgG fraction

Concentration: lot specific

Purification: Affinity Chromatography on Protein G

Conjugation: Unconjugated

Storage: Store the antibody undiluted at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.





Pecam1 Mouse Monoclonal Antibody [Clone ID: TLD-3A12] - SM272LE

Gene Name: platelet and endothelial cell adhesion molecule 1

Database Link: Entrez Gene 29583 Rat

Q3SWT0

Background: CD31, also known as platelet endothelial cell adhesion molecule 1 (PECAM1), is a type I

integral membrane glycoprotein and a member of the immunoglobulin superfamily of cell surface receptors. It is constitutively expressed on the surface of endothelial cells, and concentrated at the junction between them. It is also weakly expressed on many peripheral

lymphoid cells and platelets.

CD31 has been used to measure angiogenesis in association with tumor recurrence. Other studies have also indicated that CD31 and CD34 can be used as markers for myeloid progenitor cells and recognize different subsets of myeloid leukemia infiltrates (granular

sarcomas).

CD31 is a type 1 transmembrane protein expressed primarily on endothelial cells, platelets

and leucocytes.

Synonyms: PECAM-1, EndoCAM, GPIIA'

Note: Endotoxin Level: less than 0.01 EU/μg