

Product datasheet for SM3006PP

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD10 (MME) Mouse Monoclonal Antibody [Clone ID: MEM-78]

Product data:

Product Type: Primary Antibodies

Clone Name: MEM-78

Applications: FC

Recommended Dilution: Flow Cytometry analysis of human blood cells using 10 μl reagent / 100 μl of whole blood

or 106 cells in a suspension.

The content of a vial (1 ml) is sufficient for 100 tests.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: NALM-6 human pre-B cell line

Specificity: This antibody reacts with CD10 antigen (CALLA - Common acute lymphatic leukemia antigen),

a 100 kDa type II integral membrane protein.

Formulation: Phosphate buffered saline (PBS) solution containing 15mM sodium azide

Label: PerCP

State: Liquid Ig fraction

Label: Conjugated with Peridinin-chlorophyll-protein complex (PerCP) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for

direct use.

Conjugation: PerCP

Storage: Store undiluted at 2-8°C. DO NOT FREEZE! This products is photosensitive and should be

protected from light.

Stability: Shelf life: one year from despatch.

Gene Name: membrane metallo-endopeptidase

Database Link: Entrez Gene 4311 Human

P08473





CD10 (MME) Mouse Monoclonal Antibody [Clone ID: MEM-78] - SM3006PP

Background:

CD10 (neutral endopeptidase – NEP, common acute lymphocytic leukemia antigen – CALLA, membrane metallo-endopeptidase – MME, enkefalinase) is a 100-kDa cell surface zinc metalloprotease cleaving peptide bonds on the N-terminus of hydrophobic amino acids and inactivating multiple physiologically active peptids. CD10 is expressed on various normal cell types, including lymphoid precursor cells, germinal center B lymhocytes, and some epithelial cells, and its expression level serves as a marker for diagnostics of many carcinomas. CD10 is also a differentiation antigen for early B-lymphoid progenitors in the B-cell differentiation pathway and has a key role in regulation of growth, differentiation and signal transduction of many cellular systems.

Synonyms:

MME, EPN, Neutral endopeptidase 24.11, Neutral endopeptidase, NEP, Enkephalinase, Atriopeptidase, CALLA