

Product datasheet for **SM3011B**

CD16 (FCGR3A) Mouse Monoclonal Antibody [Clone ID: MEM-154]

Product data:

Product Type:	Primary Antibodies
Clone Name:	MEM-154
Applications:	FC
Recommended Dilution:	Flow Cytometry: 1/500 as a starting point for Indirect Immunofluorescence.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human granulocytes
Specificity:	This antibody reacts with the epitope on CD16 antigen that residing in proximity to FG loop (probably BC or C'E loop). CD16 is a low affinity receptor for aggregated IgG (FcγRIII antigen). The antibody reacts with CD16+ granulocytes.
Formulation:	PBS, pH~7.4 with 15 mM Sodium Azide as preservative. Label: Biotin State: Liquid purified Ig fraction. Label: Conjugated with -LC-NHS under optimum conditions. The reagent is free of unconjugated biotin
Concentration:	lot specific
Conjugation:	Biotin
Storage:	Store the antibody undiluted at 2 - 8 °C. DO NOT FREEZE!
Stability:	Shelf life: one year from despatch.
Gene Name:	Fc fragment of IgG receptor IIIa
Database Link:	Entrez Gene 2214 Human P08637



[View online »](#)

Background:

CD16 (FcγRIII) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human FcγRIII is expressed in two forms FcγRIII-A and -B. FcγRIII-A is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T cells. It is associated with FcεRI-γ subunit and is responsible for antibody-dependent NK cell cytotoxicity. Mast cell FcγRIII-A is associated, moreover, with FcεRI-β subunit. Besides IgG, FcεRI-A can be triggered also by oligomeric IgE. FcγRIII-B is a GPI-linked monomeric receptor expressed on neutrophils and is involved in their activation and induction of a proadhesive phenotype.

Synonyms:

FCGR3A, CD16A, FCG3, FCGR3, IGFR3, Fc-gamma RIII-alpha, Fc-gamma RIII, Fc-gamma RIIIa, FcRIII, FcRIIIa, FcR-10, IgG Fc receptor III-2