

Product datasheet for **SM3015R**

IL2 Receptor alpha (IL2RA) Mouse Monoclonal Antibody [Clone ID: MEM-181]

Product data:

Product Type:	Primary Antibodies
Clone Name:	MEM-181
Applications:	FC
Recommended Dilution:	Flow Cytometry analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	PHA-activated peripheral blood leucocytes
Specificity:	The antibody MEM-181 reacts with CD25 (Interleukin-2 receptor a chain), a 55 kDa type I transmembrane glycoprotein expressed on activated B and T lymphocytes, activated monocytes/macrophages and on CD4+ T lymphocytes (T regulatory cells); it is lost on resting B and T lymphocytes.
Formulation:	Phosphate buffered saline (PBS) containing 15 mM sodium azide Label: PE State: Liquid Ig fraction
Purification:	Size-exclusion chromatography
Conjugation:	PE
Storage:	Store the antibody undiluted at 4-8°C. DO NOT FREEZE! This antibody is photosensitive and should be protected from light. Should it contain a precipitate we recommend microcentrifugation before use.
Stability:	Shelf life: one year from despatch.
Gene Name:	interleukin 2 receptor subunit alpha
Database Link:	Entrez Gene 3559 Human P01589




[View online »](#)

Background:

CD25 (IL2Ralpha, Tac) is a ligand-binding alpha subunit of interleukin 2 receptor (IL2R). Together with beta and gamma subunit CD25 constitutes the high affinity IL2R, whereas CD25 alone serves as the low affinity IL2R. CD25 expression rapidly increases upon T cell activation. The 55 kDa CD25 molecule is enzymatically cleaved and shed from the cell surface as a soluble 45 kDa s-Tac, whose concentration in serum can be used as a marker of T cell activation. Expression of CD25 indicates the neoplastic phenotype of mast cells. Humanized anti CD25 antibodies represent a useful tool to reduce the incidence of allograft rejection as well as the severity of graft versus host reaction, and radioimmunoconjugates of anti-CD25 antibodies can be used against CD25 expressing lymphomas.

Synonyms:

Interleukin-2 receptor alpha chain, IL-2 receptor alpha subunit, IL-2-RA, IL2-RA, p55, TAC antigen

Product images:

Surface staining of PHA-stimulated human PBMC with anti-human CD25 (MEM-181) PE.