

Product datasheet for SM3081AC

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD52 Mouse Monoclonal Antibody [Clone ID: HI186]

Product data:

Product Type: Primary Antibodies

Clone Name: HI186
Applications: FC

Recommended Dilution: Flow Cytometry analysis of Human blood cells using 10 μl reagent / 100 μl of whole blood or

106 cells in a suspension.

The content of a vial (1 ml) is sufficient for 100 tests.

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal Immunogen: Human tonsil.

Specificity: This antibody reacts with Human CD52 (CAMPATH-1), a 21-28 kDa glycoprotein containing a

large N-linked carbohydrate moiety; mature CD52 molecule is actually much smaller (approx. 8-9 kDa). CD52 is expressed at high levels on lymphocytes, monocytes/macrophages and in

male reproductive tract.

Formulation: PBS

Label: APC

State: Liquid purified IgG fraction

Stabilizer: 0.2% (w/v) high-grade protease free BSA

Preservative: 15 mM Sodium Azide

Label: Conjugated with cross-linked Allophycocyanin under optimum conditions

Purification: Size-Exclusion Chromatography

Conjugation: APC

Storage: Store undiluted at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Stability: Shelf life: one year from despatch.

Gene Name: CD52 molecule



CD52 Mouse Monoclonal Antibody [Clone ID: HI186] - SM3081AC

Database Link: Entrez Gene 1043 Human

P31358

Background: CD52 (CAMPATH-1, HE5) is a highly glycosylated GPI-anchored 21-28 kDa glycopeptide which

is present at high levels on lymphocytes, macrophages, epithelial cells of male reproductive

tract and mature sperm. Its 12-amino acid beckbone carries a complex N-linked


carbohydrate moiety, which differs between sperm and leukocyte CD52, as well as the GPI anchor does. CD52 can be acquired by sperm cells from seminal plasma, where it is released by epithelial cells. Although CD52 is not an essential T-cell costimulator, its triggering results in activation of normal human T cells. CD52 is a very good target for antibody/complement-


mediated cell lysis.

Synonyms: HE5, CDw52

Protein Families: Transmembrane

Product images:





Flow cytometry analysis of CD52 in patients suffering with Acute Lymphoblastic Leukemia (anti-human CD52 (HI186); detection by Goat antimouse IgG2b PE). Fig. 1A ?? CD52 positive cALL (gate leukemic cells) Fig. 1B ?? CD52 negative pro T ALL (gate leukemic cells)