

## Product datasheet for **SM3103F**

### CD18 (ITGB2) Mouse Monoclonal Antibody [Clone ID: MEM-48]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	MEM-48
Applications:	FC, IHC, WB
Recommended Dilution:	Flow Cytometry analysis of human blood cells using 20 µl reagent / 100 µl whole blood.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Leucocytes of a patient suffering from a LGL-type leukaemia
Specificity:	The antibody recognizes an epitope involving residues 534-546 in cysteine-rich repeat 3 of the CD18 antigen (Integrin beta2 chain).
Formulation:	Phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent Label: FITC State: Liquid purified Ig fraction Label: Conjugated with Fluorescein isothiocyanate under optimum conditions. The reagent is free of unconjugated and adjusted for direct use
Conjugation:	FITC
Storage:	Store the antibody at 2 - 8 °C. DO NOT FREEZE! This product is photosensitive and should be protected from light.
Stability:	Shelf life: one year from despatch.
Gene Name:	integrin subunit beta 2
Database Link:	<a href="#">Entrez Gene 3689 Human P05107</a>



[View online »](#)

**Background:**

CD18 is a 90-95 kDa type I transmembrane protein expressed on all leukocytes. CD18, integrin  $\beta 2$  subunit, forms heterodimers with four types of CD11 molecule to constitute leukocyte ( $\beta 2$ ) integrins:  $\alpha$ L $\beta$ 2 (CD11a/CD18, LFA-1),  $\alpha$ M $\beta$ 2 (CD11b/CD18, Mac-1, CR3),  $\alpha$ X $\beta$ 2 (CD11c/CD18) and  $\alpha$ D $\beta$ 2 (CD11d/CD18). In most cases, the response mediated by the integrin is a composite of the functions of its individual subunits. These integrins are essential for proper leukocyte migration, mediating intercellular contacts. Absence of CD18 leads to leukocyte adhesion deficiency-1; severe reduction of CD18 expression leads to the development of a psoriasiform skin disease. CD18 is also a target of *Mannheimia* (*Pasteurella*) *haemolytica* leukotoxin and is sufficient to mediate leukotoxin-mediated cytolysis.

**Synonyms:**

Integrin  $\beta$ -2, MFI7




**Product images:**

Fig. 1B Western Blotting

Western Blotting analysis (non-reducing conditions) of whole cell lysate using anti-human CD18 (MEM-48). Lane 1: KG-1a human acute myeloid leukemia cell line Lane 2: HPB-ALL human peripheral blood T cell leukemia cell line



Immunohistochemistry staining of human spleen (paraffin sections) using anti-CD18 (MEM-48).



Surface staining of human peripheral blood cells with anti-human CD18 (MEM-48) PE. Cells in the granulocyte gate were used for analysis.