

Product datasheet for SM3104P

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

CD71 (TFRC) Mouse Monoclonal Antibody [Clone ID: MEM-75]

Product data:

Product Type: Primary Antibodies

Clone Name: MEM-75
Applications: FC, IF, IP

Recommended Dilution: Flow cytometry: Recommended dilution: 0.5-4 µg/ml.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Pre-B cell line NALM-6

Specificity: The antibody reacts with CD71 antigen (Transferrin Receptor), a 95 kDa type II homodimeric

transmembrane glycoprotein expressed on activated B and lymphocytes, macrophages and

erythroid precursors; it is lost on resting blood leukocytes.

The antibody does not block binding of Transferrin to the Receptor.

Formulation: Phosphate buffered saline (PBS) solution with 15 mM sodium azide

State: Purified

State: Liquid purified Ig fraction (> 98% pure by SDS-PAGE)

Concentration: lot specific

Purification: Protein-A Affinity Chromatography

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: transferrin receptor

Database Link: Entrez Gene 7037 Human

P02786





CD71 (TFRC) Mouse Monoclonal Antibody [Clone ID: MEM-75] - SM3104P


Background:

CD71 (transferrin receptor) is a type II transmembrane glycoprotein expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin and iron ions), CD71 is internalized by clathrin-mediated endocytosis. Acidification of endosomes by vesicular membrane proton pumps leads to dissociation of iron ions, whereas transferrin (apotransferrin) remains associated with CD71 and recycles to the cell surface, where it is released upon exposure to normal pH. CD71 is also involved in uptake of non-transferrin bound iron.

Synonyms:

TfR1, p90, Transferrin receptor protein 1

Product images:



Immunofluorescence staining of CD71 in human HeLa cell line using anti-CD71 (MEM-75; green). Actin cytoskeleton decorated by phalloidin (red) and cell nuclei stained with DAPI (blue).