

Product datasheet for SM311P

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MHC Class I (RT1Aa) Mouse Monoclonal Antibody [Clone ID: OX-18]

Product data:

Product Type: Primary Antibodies

Clone Name: OX-18
Applications: FC, IHC, IP

Recommended Dilution: Flow cytometry: 1:50-1:100, use 10 μl of the suggested working dilution to label 10e6 cells.

Immunohistochemistry on frozen sections: Acetone fixation recommended - the antigen is sensitive to fixation with paraformaldehyde. Not suitable on paraffin embedded sections.

ELISA. Immunoprecipitation.

Reactivity: Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Rat spleen cell glycoprotein

Specificity: This antibody recognises a monomorphic determinant of rat MHC Class I (RT-1A). However,

quantitative measurements suggest that not all of the class I molecules are recognised. MRC OX-18 has been used in immunoaffinity purification of rat MHC class I molecules (see

Ref. 1). This product is routinely tested in flow cytometry on rat splenocytes.

Formulation: PBS, pH 7.4, containing 0.09% sodium azide as preservative

State: Purified State: Liquid

Concentration: lot specific

Purification: Protein G affinity chromatography

Conjugation: Unconjugated

Storage: Store the antibody at 2-8°C for up to one month or at -20°C for longer. Avoid repeated

freezing and thawing. Should this product contain a precipitate we recommend

microcentrifugation before use.

Stability: Shelf life: one year from despatch.

Database Link: P16391







Background:

MHC Class I molecules play a central role in the immune system by presenting peptides derived from the endoplasmic reticulum lumen. MHC class I antigens are heterodimers consisting of one alpha chain (44kDa) with beta 2 microglobulin (11.5 kDa). The antigen is expressed by all somatic cells at varying levels. MHC Class I molecules are expressed on most nucleated cells where they present endogenously synthesized antigenic peptides to CD8+ T lymphocytes, which are usually cytotoxic T cells. Fibroblasts or neurons however only show a low level of antigen.