

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for SM3143P

CD41 (ITGA2B) Mouse Monoclonal Antibody [Clone ID: HIP8]

Product data:

| Product Type: | Primary Antibodies |
|-----------------------|--|
| Clone Name: | HIP8 |
| Applications: | FC, IHC |
| Recommended Dilution: | Flow Cytometry: (has been tested by flow cytometric analysis of human platelets. Can be used at less than or equal to 0.5 μg per 100 μl blood (per 10e6 cells). Immunohistochemistry (frozen sections). |
| Reactivity: | Human, Primate |
| Host: | Mouse |
| lsotype: | lgG1 |
| Clonality: | Monoclonal |
| Specificity: | The antibody reacts with alpha (a) subunit of CD41 (heavy chain; 120 kDa). CD41 is mainly expressed on platelets and megakaryocytes. The antibody blocks platelet aggregation and completely inhibits ADP-, epinephrine-, and collagen-induced platelet activation, and partially inhibits ristocetin- and thrombin-induced platelet activation. It is useful in the morphological and physiological studies of platelets and megakaryocytes. |
| Formulation: | Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4 State: Purified State: Liquid Ig fraction |
| Concentration: | lot specific |
| Purification: | Protein-G affinity chromatography; purity: > 95% (by SDS-PAGE) |
| Conjugation: | Unconjugated |
| Storage: | Store the antibody at 2 - 8 °C up to one month or (in aliquots) at -20 °C for longer. Avoid repeated freezing and thawing. |
| Stability: | Shelf life: one year from despatch. |
| Gene Name: | integrin subunit alpha 2b |
| Database Link: | <u>Entrez Gene 3674 Human</u> <u>P08514</u> |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

CD41 (ITGA2B) Mouse Monoclonal Antibody [Clone ID: HIP8] – SM3143P Background: CD41 (platelet glycoprotein IIb) is composed of two subunits (120 kDa a, a and 23 kDa b, b) that interact with CD61 in the presence of calcium to form a functional adhesive protein receptor. Upon blood vessel damage, this receptor binds to a variety of proteins including von Willebrand factor, fibrinogen, fibronectin and vitronectin. CD41 is mainly expressed on megakaryocyte-platelet lineage, but generally belongs to the antigens that are expressed during early stages of hematopoietic differentiation.

Synonyms:

Integrin alpha-IIb, GP2B, ITGAB, GPalpha IIb

Product images:



Surface staining of human peripheral blood cells with anti-CD41a (HIP8) purified, GAM-APC.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US