

Product datasheet for SM3153B

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD44 Mouse Monoclonal Antibody [Clone ID: MEM-85]

Product data:

Product Type: Primary Antibodies

Clone Name: MEM-85

Applications: FC

Recommended Dilution: Indirect immunofluorescence analysis by **Flow Cytometry**: 1:1000 starting point.

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Leukocytes of a patient suffering from LGL Type Leukaemia

Specificity: The antibody MEM-85 reacts with both cell surface-expressed and soluble form of CD44

antigen (Phagocyte glycoprotein 1), a 80-95 kDa transmembrane glycoprotein (hyaladherin family) present on the most of cells and tissues (leukocytes, endothelial cells, mesenchymal

cells, etc.); it is negative on platelets and hepatocytes.

Formulation: Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4

Label: Biotin

State: Liquid purified Ig fraction

Label: Conjugated with Biotin-LC-NHS under optimum conditions. The reagent is free of

unconjugated biotin.

Concentration: lot specific
Conjugation: Biotin

Storage: Store the antibody undiluted at 2 - 8 °C. DO NOT FREEZE!

Stability: Shelf life: one year from despatch.

Gene Name: CD44 molecule (Indian blood group)

Database Link: Entrez Gene 960 Human

P16070





Background:

CD44 is a transmembrane glycoprotein expressed on the surface of most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculation and homing, it can thus serve e.g. as a modulator of macrophage recruitment in response to pathogen. Although CD44 functions are essential for physiological activities of normal cells, elevated CD44 expression correlates with poor prognosis in many carcinomas, facilitating tumour growth and metastasis, antiapoptosis and directional motility of cancer cells.


Synonyms: LHR, MDU2, MDU3, MIC4, CDw44, Epican, ECMR-III, HUTCH-I, Heparan sulfate proteoglycan,

Hermes antigen, Hyaluronate receptor, PGP-1

Note: HLDA IV; WS Code NL 706

HLDA VI; WS Code AS Ref.15

Product images:



Surface staining of human peripheral blood leukocytes with anti-human CD44 (MEM-85) biotin.