

Product datasheet for SM502R

CD44 Mouse Monoclonal Antibody [Clone ID: CVS18]

Product data:

Product Type: Primary Antibodies

Clone Name: CVS18

Applications: FC

Recommended Dilution: Flow cytometry: Neat, use 10μl to label 106 cells in 100μl.

Reactivity: Equine
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Equine leucocytes

Specificity: Clone CVS18 antibody detects equine CD44 and may be used as a pan equine leucocyte

marker.

Formulation: 0.09% Sodium azide, 1% BSA and 5% Sucrose

Label: PE State: Purified

State: Lyophilised purified IgG

Label: R. Phycoerythrin

Reconstitution Method: Restore with 1.0 ml distilled water.

Purification: Affinity chromatography on Protein G

Conjugation: PE

Storage: Prior to and following reconstitution store the antibody at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Stability: Shelf life: one year from despatch.

Database Link: Q05078



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



CD44 Mouse Monoclonal Antibody [Clone ID: CVS18] - SM502R

Background:

CD44 is a type 1 transmembrane glycoprotein also known as Phagocytic Glycoprotein 1 (pgp 1) and HCAM. CD44 is the receptor for hyaluronate and exists as a large number of different isoforms due to alternative RNA splicing. The major isoform expressed on lymphocytes, myeloid cells, and erythrocytes is a glycosylated type 1 transmembrane protein. Other isoforms contain glycosaminoglycans and are expressed on hematopoietic and non hematopoietic cells. CD44 is involved in adhesion of leukocytes to endothelial cells, stromal cells, and the extracellular matrix.

Synonyms:

LHR, MDU2, MDU3, MIC4, CDw44, Epican, ECMR-III, HUTCH-I, Heparan sulfate proteoglycan, Hermes antigen, Hyaluronate receptor, PGP-1