

Product datasheet for SM6013S

MIF Mouse Monoclonal Antibody [Clone ID: 4E4]

Product data:

Product Type: Primary Antibodies Clone Name: 4F4 **Applications:** ELISA, FC, IF, IHC, WB Recommended Dilution: ELISA. Western blot: 1/500-1/2,000. Recommended starting dilution is 1/1,000. Flow Cytometry. Immunofluorescence/Immnunocytochemistry. Immunohistochemistry on Paraffin Sections: 5 µg/ml. Human **Reactivity:** Host: Mouse lgG1 Isotype: **Clonality:** Monoclonal Immunogen: Recombinant Human MIF (1-115 aa) purified from E. coli. The antibody recognizes Human MIF. Specificity: Other species not tested. Formulation: PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol State: Purified State: Liquid purified Ig fraction Concentration: lot specific **Purification:** Affinity Chromatography on Protein G **Conjugation:** Unconjugated Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Stability: Shelf life: one year from despatch. Gene Name: macrophage migration inhibitory factor (glycosylation-inhibiting factor) Database Link: Entrez Gene 4282 Human P14174




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| | MIF Mouse Monoclonal Antibody [Clone ID: 4E4] – SM6013S |
|-------------|---|
| Background: | The cytokine Macrophage migration inhibitory factor (MIF) has been identified to be secreted by the pituitary gland and the monocyte/macrophage and to play an important role in endotoxic shock. MIF has the unique property of being released from macrophages and T- cells in response to physiological concentrations of glucocorticoids. The secretion of MIF is tightly regulated and decreases at high, anti-inflammatory steroid concentration. |
| Synonyms: | Macrophage migration inhibitory factor, Phenylpyruvate tautomerase, Glycosylation- inhibiting factor, GLIF, MMIF |

Product images:



Western blot analysis: The extract of HL-60 was resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-Human MIF antibody (1/1,000). Protein wasvisualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.



[SM6013] MIF antibody staining of Formalin-Fixed, Paraffin-Embedded Human Breast.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US