

Product datasheet for SM6024S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GST-Tag Mouse Monoclonal Antibody [Clone ID: 1E5]

Product data:

Product Type: Primary Antibodies

Clone Name: 1E5

Applications: ELISA, IF, WB

Recommended Dilution: ELISA.

Western blot (1/500-1/2,000), recommended starting dilution is 1/1,000.

Immunofluorescence/Immnunocytochemistry.

Reactivity: Schistosoma japonicum

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Recombinant Glutathione S transferase (GST) purified from *E. coli*

Specificity: This antibody reacting specifically with GST will be useful in various immunotechniques to

identify the expression of a GST fusion protein in bacteria, bacterial lysates or cells and

tissues transfected with a GST fusion protein expressing vectors.

Formulation: PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol

State: Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Affinity Chromatography on Protein G

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Database Link: P08515






Background:

Glutathione S transferases (GSTs) are a family of enzymes that play an important role in detoxification by catalyzing the conjugation of many hydrophobic and electrophilic compounds with reduced glutathione. GST proteins can be purified by immobilized glutathione affinity chromatography. Many recombinant proteins have been engineered with GST tags to facilitate the detection, isolation and purification of these proteins.

Synonyms:

Glutathione S-transferase Tag, GST26-Tag

Product images:



Western blot analysis: Recombinant GST (28 kDa) and GST-fusion protein (61 kDa) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-GST antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugat