

Product datasheet for **SP2088P**

Ccl24 Goat Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, FN, WB
Recommended Dilution:	Neutralisation: To yield one-half maximal inhibition [ND50] of the biological activity of mEotaxin-2 (100 ng/ml), a concentration of 7.0-8.0 µg/ml of this antibody is required. Sandwich ELISA: To detect mEotaxin-2 (using 100 µl/well antibody solution) a concentration of 0.5-2.0 µg/ml of this antibody is required. In conjunction with Biotinylated Anti-Murine Eotaxin-2 as a detection antibody, it allows the detection of at least 0.2-0.4 ng/well of recombinant mEotaxin-2. Indirect ELISA: To detect mEotaxin-2 (using 100 µl/well antibody solution) a concentration of 0.5-2.0 µg/ml of this antibody is required. In conjunction with compatible secondary reagents, it allows the detection of at least 0.2-0.4 ng/well of recombinant mEotaxin-2. Western blot: To detect mEotaxin-2 this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mEotaxin-2 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Mouse
Host:	Goat
Clonality:	Polyclonal
Immunogen:	Recombinant mouse Eotaxin-2
Specificity:	This antibody detects Eotaxin-2.
Formulation:	PBS, pH 7.2 State: Aff - Purified State: Sterile filtered lyophilized Ig fraction
Reconstitution Method:	Centrifuge vial prior to opening. Restore in sterile water to a concentration of 0.1 - 1.0 mg/ml.
Purification:	Immunoaffinity chromatography
Conjugation:	Unconjugated
Storage:	Store lyophilized at 2-8°C for 6 months or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing.



[View online »](#)

Stability:	Shelf life: one year from despatch.
Gene Name:	chemokine (C-C motif) ligand 24
Database Link:	Entrez Gene 56221 Mouse Q9JKC0
Background:	<p>Eotaxin 2 is a member of the CC chemokine family, based on the presence of the CC motif and homology with other known CC chemokines. Eotaxin 2 cDNA encodes a 119 amino acid residue precursor protein with a 26 amino acid residue signal peptide that is cleaved to generate a mature protein predicted to contain 93 amino acid residues with an N glycosylation site. Mature human eotaxin 2 has a predicted molecular mass of approximately 10.6 kDa. Compared to other CC chemokines, eotaxin 2 exhibits 40 %, 42 %, and 39 % amino acid identity to MCP3, MIP1 alpha, and eotaxin, respectively. Human CC chemokine eotaxin 2 maps to chromosome 7q11.23.</p> <p>Both eotaxin and eotaxin 2 activate and attract eosinophils and basophils. A receptor for human eotaxin has been identified and found to be the third numbered receptor in the C-C chemokine subfamily of receptors (CCR3. On eosinophils, the effects of eotaxin 2 is inhibited by an CCR3 antibody and cross-desensitized by eotaxin and MCP4, suggesting that all three CC chemokines act through CCR3. Eotaxin 2 mRNA is weakly expressed in activated monocytes and T lymphocytes. Recombinant eotaxin 2 induces chemotaxis of eosinophils, basophils, and resting T lymphocytes but not monocytes and activated T lymphocytes. Eotaxin 2 inhibits colony formation in myeloid progenitor cells.</p>
Synonyms:	C-C motif chemokine 24, Small-inducible cytokine A24, MPIF2, MPIF-2, SCYA24, CK-beta-6