

## Product datasheet for **SR300489**

### **BYSL Human siRNA Oligo Duplex (Locus ID 705)**

#### **Product data:**

<b>Product Type:</b>	siRNA Oligo Duplexes
<b>Purity:</b>	HPLC purified
<b>Quality Control:</b>	Tested by ESI-MS
<b>Sequences:</b>	Available with shipment
<b>Stability:</b>	One year from date of shipment when stored at -20°C.
<b># of transfections:</b>	Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).
<b>Note:</b>	Single siRNA duplex (10nmol) can be ordered.
<b>RefSeq:</b>	<a href="#">NM_004053</a>
<b>UniProt ID:</b>	<a href="#">Q13895</a>
<b>Synonyms:</b>	BYSTIN; Enp1
<b>Components:</b>	BYSL (Human) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 705) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNase free siRNA Duplex Resuspension Buffer - 2 ml
<b>Summary:</b>	<p>Bystin is expressed as a 2-kb major transcript and a 3.6-kb minor transcript in SNG-M cells and in human trophoblastic teratocarcinoma HT-H cells. Protein binding assays determined that bystin binds directly to trophinin and tastin, and that binding is enhanced when cytokeratins 8 and 18 are present. Immunocytochemistry of HT-H cells showed that bystin colocalizes with trophinin, tastin, and the cytokeratins, suggesting that these molecules form a complex in trophoblast cells at the time of implantation. Using immunohistochemistry it was determined that trophinin and bystin are found in the placenta from the sixth week of pregnancy. Both proteins were localized in the cytoplasm of the syncytiotrophoblast in the chorionic villi and in endometrial decidual cells at the uteroplacental interface. After week 10, the levels of trophinin, tastin, and bystin decreased and then disappeared from placental villi. [provided by RefSeq, Jul 2008]</p>



[View online »](#)

**Performance  
Guaranteed:**

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at [techsupport@origene.com](mailto:techsupport@origene.com). Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).