

Product datasheet for SR303924

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PEX19 Human siRNA Oligo Duplex (Locus ID 5824)

Product data:

Product Type: siRNA Oligo Duplexes

Purity: HPLC purified

Quality Control: Tested by ESI-MS

Sequences: Available with shipment

Stability: One year from date of shipment when stored at -20°C.

of transfections: Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final

conc. 10 nM).

Note: Single siRNA duplex (10nmol) can be ordered.

RefSeq: NM 001131039, NM 001193644, NM 002857, NR 036492, NR 036493

UniProt ID: P40855

Synonyms: D1S2223E; FLJ55296; HK33; housekeeping gene, 33kD; OTTHUMP00000031848; peroxisomal

biogenesis factor 19; peroxisomal farnesylated protein; PMP1; PMPI; PXF; PXMP1

Components: PEX19 (Human) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 5824)

Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol

Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 ml

Summary: This gene is necessary for early peroxisomal biogenesis. It acts both as a cytosolic chaperone

and as an import receptor for peroxisomal membrane proteins (PMPs). Peroxins (PEXs) are proteins that are essential for the assembly of functional peroxisomes. The peroxisome biogenesis disorders (PBDs) are a group of genetically heterogeneous autosomal recessive, lethal diseases characterized by multiple defects in peroxisome function. These disorders have at least 14 complementation groups, with more than one phenotype being observed for some complementation groups. Although the clinical features of PBD patients vary, cells from all PBD patients exhibit a defect in the import of one or more classes of peroxisomal matrix proteins into the organelle. Defects in this gene are a cause of Zellweger syndrome (ZWS), as well as peroxisome biogenesis disorder complementation group 14 (PBD-CG14), which is also known as PBD-CGJ. Alternative splicing results in multiple transcript variants. [provided by

RefSeq, Aug 2010]







Performance Guaranteed:

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).