

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for SR305846

## PAG3 (ASAP2) Human siRNA Oligo Duplex (Locus ID 8853)

## **Product data:**

Product Type:	siRNA Oligo Duplexes
Purity:	HPLC purified
Quality Control:	Tested by ESI-MS
Sequences:	Available with shipment
Stability:	One year from date of shipment when stored at -20°C.
# of transfections:	Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).
Note:	Single siRNA duplex (10nmol) can be ordered.
RefSeq:	<u>NM 001135191, NM 003887</u>
UniProt ID:	<u>O43150</u>
Synonyms:	AMAP2; CENTB3; DDEF2; PAG3; PAP; Pap-alpha; SHAG1
Components:	ASAP2 (Human) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 8853) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 ml
Summary:	<ul> <li>This gene encodes a multidomain protein containing an N-terminal alpha-helical region with a coiled-coil motif, followed by a pleckstrin homology (PH) domain, an Arf-GAP domain, an ankyrin homology region, a proline-rich region, and a C-terminal Src homology 3 (SH3) domain. The protein localizes in the Golgi apparatus and at the plasma membrane, where it colocalizes with protein tyrosine kinase 2-beta (PYK2). The encoded protein forms a stable complex with PYK2 in vivo. This interaction appears to be mediated by binding of its SH3 domain to the C-terminal proline-rich domain of PYK2. The encoded protein is tyrosine phosphorylated by activated PYK2. It has catalytic activity for class I and II ArfGAPs in vitro, and can bind the class III Arf ARF6 without immediate GAP activity. The encoded protein is believed to function as an ARF GAP that controls ARF-mediated vesicle budding when recruited to Golgi membranes. In addition, it functions as a substrate and downstream target for PYK2 and SRC, a pathway that may be involved in the regulation of vesicular transport. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2008]</li> </ul>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2021 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	PAG3 (ASAP2) Human siRNA Oligo Duplex (Locus ID 8853) – SR305846
Performance	OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will
Guaranteed:	provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.
	For non-conforming siRNA, requests for replacement product must be made within ninety

(90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2021 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US