

## **Product datasheet for SR310535**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RFWD3 Human siRNA Oligo Duplex (Locus ID 55159)

#### **Product data:**

**Product Type:** siRNA Oligo Duplexes

Purity: HPLC purified

**Quality Control:** Tested by ESI-MS

Sequences: Available with shipment

**Stability:** One year from date of shipment when stored at -20°C.

# of transfections: Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final

conc. 10 nM).

**Note:** Single siRNA duplex (10nmol) can be ordered.

**RefSeq:** NM 018124, NM 001370536, NM 001370534, NM 001370535, NM 001370539,

NM 001370540, NM 001370543, NM 001370537, NM 001370542

UniProt ID: Q6PCD5

Synonyms: FANCW; RNF201

Components: RFWD3 (Human) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 55159)

Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol

Included - SR30005, RNAse free siRNA Duplex Resuspension Buffer - 2 ml

**Summary:** E3 ubiquitin-protein ligase required for the repair of DNA interstrand cross-links (ICL) in

response to DNA damage (PubMed:21504906, PubMed:21558276, PubMed:26474068, PubMed:28575657, PubMed:28575658). Plays a key role in RPA-mediated DNA damage

signaling and repair (PubMed:21504906, PubMed:21558276, PubMed:26474068,

PubMed:28575657, PubMed:28575658, PubMed:28691929). Acts by mediating ubiquitination of the RPA complex (RPA1, RPA2 and RPA3 subunits) and RAD51 at stalled replication forks, leading to remove them from DNA damage sites and promote homologous recombination (PubMed:26474068, PubMed:28575657, PubMed:28575658). Also mediates the ubiquitination

of p53/TP53 in the late response to DNA damage, and acts as a positive regulator of

p53/TP53 stability, thereby regulating the G1/S DNA damage checkpoint (PubMed:20173098). May act by catalyzing the formation of short polyubiquitin chains on p53/TP53 that are not targeted to the proteasome (PubMed:20173098). In response to ionizing radiation, interacts with MDM2 and enhances p53/TP53 ubiquitination, possibly by restricting MDM2 from

extending polyubiquitin chains on ubiquitinated p53/TP53 (PubMed:20173098).

[UniProtKB/Swiss-Prot Function]







# Performance Guaranteed:

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).