

Product datasheet for **SR310855**

DSU (MREG) Human siRNA Oligo Duplex (Locus ID 55686)

Product data:

Product Type:	siRNA Oligo Duplexes
Purity:	HPLC purified
Quality Control:	Tested by ESI-MS
Sequences:	Available with shipment
Stability:	One year from date of shipment when stored at -20°C.
# of transfections:	Approximately 330 transfections/2nmol in 24-well plate under optimized conditions (final conc. 10 nM).
Note:	Single siRNA duplex (10nmol) can be ordered.
RefSeq:	<u>NM_018000</u>
UniProt ID:	<u>Q8N565</u>
Synonyms:	DSU; WDT2
Components:	MREG (Human) - 3 unique 27mer siRNA duplexes - 2 nmol each (Locus ID 55686) Included - SR30004, Trilencer-27 Universal Scrambled Negative Control siRNA Duplex - 2 nmol Included - SR30005, RNase free siRNA Duplex Resuspension Buffer - 2 ml



[View online »](#)

Summary:

Probably functions as cargo-recognition protein that couples cytoplasmic vesicles to the transport machinery. Plays a role in hair pigmentation, a process that involves shedding of melanosome-containing vesicles from melanocytes, followed by phagocytosis of the melanosome-containing vesicles by keratinocytes. Functions on melanosomes as receptor for RILP and the complex formed by RILP and DCTN1, and thereby contributes to retrograde melanosome transport from the cell periphery to the center. Overexpression causes accumulation of late endosomes and/or lysosomes at the microtubule organising center (MTOC) at the center of the cell. Probably binds cholesterol and requires the presence of cholesterol in membranes to function in microtubule-mediated retrograde organelle transport. Binds phosphatidylinositol 3-phosphate, phosphatidylinositol 4-phosphate, phosphatidylinositol 5-phosphate and phosphatidylinositol 3,5-bisphosphate, but not phosphatidylinositol 3,4-bisphosphate or phosphatidylinositol 4,5-bisphosphate (By similarity). Required for normal phagosome clearing and normal activation of lysosomal enzymes in lysosomes from retinal pigment epithelium cells (PubMed:19240024). Required for normal degradation of the lipofuscin component N-retinylidene-N-retinylethanolamine (A2E) in the eye. May function in membrane fusion and regulate the biogenesis of disk membranes of photoreceptor rod cells (By similarity).[UniProtKB/Swiss-Prot Function]

Performance Guaranteed:

OriGene guarantees that at least two of the three Dicer-Substrate duplexes in the kit will provide at least 70% or more knockdown of the target mRNA when used at 10 nM concentration by quantitative RT-PCR when the TYE-563 fluorescent transfection control duplex (cat# SR30002) indicates that >90% of the cells have been transfected and the HPRT positive control (cat# SR30003) provides 90% knockdown efficiency.

For non-conforming siRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the siRNA kit. To arrange for a free replacement with newly designed duplexes, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled siRNA control (quantitative RT-PCR data required).